

Interferometry with Two Pairs of Spin Correlated Photons

Mladen Pavičić

*Institut für Theoretische Physik, TU Berlin, Hardenbergstraße 36, D-10623 Berlin 12, Germany
 Atominstytut der Österreichischen Universitäten, Schüttelstraße 115, A-1020 Wien, Austria
 and Department of Mathematics, University of Zagreb, GF, Kaciceva 26, HR-41001 Zagreb, Croatia**

Johann Summhammer

*Atominstytut der Österreichischen Universitäten, Schüttelstraße 115, A-1020 Wien, Austria
 (Received 5 November 1993)*

We propose a new experiment employing two independent sources of spin correlated photon pairs. Two photons from different unpolarized sources each pass through a polarizer to a detector. Although their trajectories never mix or cross, they exhibit fourth-order-interference-like correlations when the other two photons interfere on a beam splitter even when the latter two do not pass any polarizers at all. A wave packet calculation shows that the experiment permits a very discriminatory test of hidden variable theories.

PACS numbers: 03.65.Bz, 42.50.Wm

Higher order interference effects have been much investigated because their nonlocal nature provides a powerful tool for testing hidden variables theories [1–15]. In recent test Ou and Mandel [7] used the polarization correlation of signal and idler photon of down-converted light and found a violation of Bell's inequality by about 6 standard deviations. However, their detectors were not sufficiently efficient. The idea was extended to two-particle interferometry in a proposal by Horne, Shimony, and Zeilinger [11]. In another recent experiment Wang, Zou, and Mandel [12] tested the so-called de Broglie-Bohm pilot wave theory. The result was negative but the setup was recognized to lack generality by Holland and Vigier [13] and by Wang, Zou, and Mandel [14]. Thus excluding realistic nonlocality remains a challenge [16].

In this paper we propose an experiment which should be the first realization of the fourth-order interference of randomly prepared independent photons correlated in polarization and coming from independent sources. The experiment is based on a newly discovered interference effect of the fourth-order on a beam splitter [17]. The essential new element of the experiment is that it puts together systems which were not in any way influenced by preparation, makes them interact, and then allows us inferring overall polarization (spin) correlations—from the assumed quantum mechanical description of the unknown initial states—by simultaneous measurement of four photons separated in space. Particular polarization (spin) correlations, unexpectedly found between photons which did not in any way directly interact and on the distant pairs of which polarization has not been measured at all, are expected to be confirmed by a future experiment. Such experiments might eventually disprove any realistic hidden variable theory.

A schematic representation of the experiment is shown in Fig. 1. Two independent sources, S_I and S_{II} , both simultaneously emit two photons correlated in polarization

to the left and right. On the left photons we measure polarizations by the polarization filters P_1 and P_2 and on the right photons by P_3 and P_4 . Because of the beam splitter the paths leading through P_3 and P_4 are available to the right photons from both sources S_I and source S_{II} . The resulting fourth-order interference will manifest itself in the probability of quadruple coincidence counts in detectors D_1 , D_2 , D_3 , and D_4 .

The sources can be atoms exhibiting cascade emission. (Down-conversion is not possible because it gives polarized photons.) The atoms of the two sources could be pumped to an upper level by two independent lasers [9]. This level would decay by emitting two photons correlated in polarization [18]. The independence of the two sources can be assured by slight differences in central frequency and drift of the two pump lasers. Hence there should be no second-order interference at detectors D_3 and D_4 , which could also be suppressed when the size of the sources exceeds the coherence length of the emitted photons. Interference of fourth order would still occur, because its relevant coherence time is given by the inverse of the frequency difference of the two correlated photons [1–4].

The state of the four photons immediately after leaving the sources is described by the product of two entangled




FIG. 1. Layout of the proposed experiment.

states:

$$|\Psi\rangle = \frac{1}{\sqrt{2}} (|1_x\rangle_1 |1_x\rangle_3 + |1_y\rangle_1 |1_y\rangle_3) \otimes \frac{1}{\sqrt{2}} (|1_x\rangle_2 |1_x\rangle_4 + |1_y\rangle_2 |1_y\rangle_4). \quad (1)$$

Here, $|x\rangle$ and $|y\rangle$ denote the mutually orthogonal photon states. So, e.g., $|1_x\rangle_1$ means the state of photon 1 leaving the source S_1 to the left polarized in direction x . In the following we use the annihilation operator formalism, often employed in quantum optical analysis, e.g., by Paul [3], Mandel's group [5,6,9], and Campos, Saleh, and Teich [10]. The operator describing the polarization at $P1$ oriented along the x axis and the subsequent detection at $D1$ acts as follows: $\hat{a}_{1x}|1_x\rangle_1 = |0_x\rangle_1$, $\hat{a}_{1x}^\dagger|0_x\rangle_1 = |1_x\rangle_1$, $\hat{a}_{1x}|0_x\rangle_1 = 0$, etc. [3]. When $P1$ is oriented at some angle θ_1 , polarization and detection are represented by $\hat{a}_1 = \hat{a}_{1x} \cos \theta_1 + \hat{a}_{1y} \sin \theta_1$. The phase the photon accumulates between the source S_1 and the detector $D1$ adds the factor $e^{i\omega_1(r_1/c + t_0^1 - t_1)}$, where ω_1 is the frequency of photon 1, r_1 is the path length from S_1 to $D1$, c is the velocity of light, t_0^1 is the time of emission of a pair of photons at S_1 , and t_1 is the time of detection at $D1$. Hence the annihilation of a photon at detector $D1$ means application of the operator $\hat{E}_1 = (\hat{a}_{1x} \cos \theta_1 + \hat{a}_{1y} \sin \theta_1) e^{i\omega_1(r_1/c + t_0^1 - t_1)}$ onto the initial state of Eq. (1). Similarly, detection of photon 2 at $D2$ means application

of $\hat{E}_2 = (\hat{a}_{2x} \cos \theta_2 + \hat{a}_{2y} \sin \theta_2) e^{i\omega_2(r_2/c + t_0^{II} - t_2)}$, where the symbols are defined by analogy. On the right side of the sources, a detection at $D3$ can be caused by photon 3 emitted by source S_1 or by photon 4 emitted by source S_{II} . The beam splitter BS may have polarization dependent transmission and reflection coefficients, denoted by T_x, T_y , and R_x, R_y , respectively. The angle of the polarizer $P3$ is given by θ_3 . Hence, we obtain [19]

$$\hat{E}_3 = (\hat{a}_{3x} \sqrt{T_x} \cos \theta_3 + \hat{a}_{3y} \sqrt{T_y} \sin \theta_3) e^{i\omega_3[(r_{II} + r_3/c) + t_0^{II} - t_3]} + i(\hat{a}_{3x} \sqrt{R_x} \cos \theta_3 + \hat{a}_{3y} \sqrt{R_y} \sin \theta_3) e^{i\omega_3[(r_1 + r_3/c) + t_0^1 - t_3]}.$$

With similar arguments the interactions leading to registration of a photon at detector $D4$ are given by the operator

$$\hat{E}_4 = (\hat{a}_{4x} \sqrt{T_x} \cos \theta_4 + \hat{a}_{4y} \sqrt{T_y} \sin \theta_4) e^{i\omega_3[(r_{II} + r_4/c) + t_0^{II} - t_4]} + i(\hat{a}_{4x} \sqrt{R_x} \cos \theta_4 + \hat{a}_{4y} \sqrt{R_y} \sin \theta_4) e^{i\omega_3[(r_1 + r_4/c) + t_0^1 - t_4]}.$$

Here, r_1 and r_{II} denote the distance from the respective source to the beam splitter, r_3 denotes the distance from the beam splitter to detector $D3$, t_3 is the time of detection at $D3$, and ω_3 is the frequency of photon 3. The symbols r_4 , t_4 , and ω_4 are defined analogously. The evolution of the initial state through interaction with the whole setup including detection of one photon in each detector is then given by

$$\hat{E}_4 \hat{E}_3 \hat{E}_2 \hat{E}_1 |\Psi\rangle = e^{i[\omega_1(\frac{r_1}{c} + t_0^1 - t_1) + \omega_2(\frac{r_2}{c} + t_0^{II} - t_2) + \omega_3(\frac{r_3}{c} + t_0^1) + \omega_4(\frac{r_4}{c} + t_0^{II})]} \times \{T_{14}T_{23} e^{i[\omega_3(\frac{r_3}{c} - t_4) + \omega_4(\frac{r_3}{c} - t_3)]} - R_{24}R_{13} e^{i[\omega_3(\frac{r_3}{c} - t_3) + \omega_4(\frac{r_3}{c} - t_4)]}\} |0\rangle, \quad (2)$$

where

$$T_{ij} = \sqrt{T_x} \cos \theta_i \cos \theta_j + \sqrt{T_y} \sin \theta_i \sin \theta_j, \\ R_{ij} = \sqrt{R_x} \cos \theta_i \cos \theta_j + \sqrt{R_y} \sin \theta_i \sin \theta_j. \quad (3)$$

The squared modulus of this result gives the probability of having one photon arrive in each detector as a function of the angles of the polarizers. Note that the interference term, which is the real part of the product of the two terms in brackets, contains only the detection times t_3 and t_4 at detectors $D3$ and $D4$, respectively. Hence we would expect the fourth-order interference to occur only in the coincidence counts of those two detectors. However, assuming for simplicity $r_1 = r_{II}$, $r_3 = r_4$, $\omega_3 = \omega_4$, and $T_x = T_y = R_x = R_y = 1/2$, we get for the coincidence probability

$$P(\theta_1, \theta_2, \theta_3, \theta_4) = \langle \Psi | \hat{E}_1^\dagger \hat{E}_2^\dagger \hat{E}_3^\dagger \hat{E}_4^\dagger \hat{E}_4 \hat{E}_3 \hat{E}_2 \hat{E}_1 | \Psi \rangle \\ = \frac{1}{16} \sin^2(\theta_1 - \theta_2) \sin^2(\theta_3 - \theta_4). \quad (4)$$

The correlations thus exist between the polarizations on the right side *and between those on the left side*. This means that the two photons going to the left can be forced into a nonlocal polarization correlation although they are emitted from two independent sources and nowhere share

a common trajectory. Moreover, the correlation does not depend on the frequencies of the two photons, nor is there any condition as to the permissible time interval $|t_1 - t_2|$ between their detections. This can be seen in Eq. (2) where the relevant parameters ($\omega_1, \omega_2, t_1, t_2, r_1, r_2$) only enter in the overall phase factor in contrast to photons 3 and 4, which must be detected within a time interval shorter than the beating period $|\omega_3 - \omega_4|^{-1}$.

It is now interesting to see that the nonlocal polarization correlation between the photons on one side persists even when the polarizers on the other side are removed. Without the polarizers $P1$ and $P2$ of the left side, we have to sum over the probabilities of the four possible orthogonal settings of these polarizers and obtain

$$P(\infty, \infty, \theta_3, \theta_4) = \frac{1}{8} \left\{ 1 - \cos^2(\theta_3 - \theta_4) \cos \left[(\omega_3 - \omega_4) \times \left(\frac{r_4 - r_3}{c} + t_3 - t_4 \right) \right] \right\}, \quad (5)$$

where we set $r_1 = r_{II}$ and $T_x = T_y = R_x = R_y = 1/2$, but otherwise put no restrictions on the frequencies of the photons, the path lengths to the detectors, or the detection times. Equation (5) expresses the kind of nonlocal correlations from independent sources proposed by Yurke and Stoler [20].

On the other hand, if we remove the polarizers of the right side, $P3$ and $P4$, we find

$$P(\theta_1, \theta_2, \infty, \infty) = \frac{1}{8} \left\{ 1 - \cos^2(\theta_1 - \theta_2) \cos \left[(\omega_3 - \omega_4) \times \left(\frac{r_4 - r_3}{c} + t_3 - t_4 \right) \right] \right\}. \quad (6)$$

We see here explicitly, that the two photons on the left side can show a nonlocal correlation only if the time interval between detections on the right side, $|t_3 - t_4|$, is kept below the beating frequency of the photons on the right side. This condition becomes trivial when we have $\omega_3 = \omega_4$. Then the above expression suggests that there should be a time independent nonlocal polarization correlation between photons 1 and 2, although no polarization is measured on photons 3 and 4. At first glance this appears to be equivalent to having two independent one-particle sources emitting unpolarized photons of different frequencies, which nowhere meet before their detection, and which yet are supposed to show nonlocal polarization correlations that are constant in time, and we would expect no polarization correlation in such a situation. But also the result of Eq. (5) is surprising, because it is what Ou and Mandel obtained [7,8] for the interference of the two orthogonally polarized signal and idler beams of down-converted light, while we are dealing with originally unpolarized beams.

In order to see whether the nonlocal correlations of Eqs. (4) and (5) can be used to test Bell's inequality we turn to a more realistic description by means of wave packets. Each photon is represented by a Gaussian amplitude distribution of energies. For the sake of simplicity all four photons shall have the same width of the energy distribution and thus the same coherence time T . The probability amplitude that photon i has frequency ω_i when its central frequency is ω_{i0} is given by

$$f(\omega_i, \omega_{i0}, T) = \frac{T^{1/2}}{\pi^{1/4}} e^{-(\omega_i - \omega_{i0})^2 T^2/2} \quad \text{with } i = 1, \dots, 4,$$

where we normalized $|f_i|^2$ to 1. The final state of Eq. (2) must be multiplied with these four functions, and integrations must be made over the frequencies $\omega_i, i = 1, \dots, 4$. This models a photon pair as two wave packets fully overlapping at the source at the time of emission and then moving apart. Now Eqs. (4) and (5) turn into

$$P(\infty, \infty, \theta_3, \theta_4) = \frac{F}{T^4} \{ \cosh(\tau_s \tau_{34}/T^2) - \cos^2(\theta_3 - \theta_4) \cos[(\omega_3 - \omega_4)\tau_{34}] \},$$

$$P(\theta_1, \theta_2, \infty, \infty) = \frac{F}{T^4} \{ \cosh(\tau_s \tau_{34}/T^2) - \cos^2(\theta_1 - \theta_2) \cos[(\omega_3 - \omega_4)\tau_{34}] \}, \quad (7)$$

where we set $r_3 = r_4$ and defined $\tau_s \equiv t_0^I - t_0^{II}$ and $\tau_{34} \equiv t_3 - t_4$. We are now dealing with probability densities for a quadruple detection at time points t_1, t_2, t_3 , and t_4 . The damping term F contains the detection times t_1 and t_2 and other experimental parameters. It expresses how

well the wave packets are centered at the various detectors at the respective detection times. Note that in both cases the polarization correlations persist but are reduced in visibility, which is given by $v = [2 \cosh(\tau_s \tau_{34}/T^2) - 1]^{-1}$ when $(\omega_{30} - \omega_{40})\tau_{34} = 0$. For a violation of Bell's inequality, and hence for a possible exclusion of hidden variable theories, v must be larger than $2^{-1/2}$ implying the product $\tau_{34}\tau_s$ must be less than $0.663T^2$. However, the time interval between the emissions at the two sources, τ_s , is not a directly measurable quantity, but must be inferred from the detection times. Therefore τ_s cannot be known better than to about T . Hence $\tau_{34} \ll T$ is a necessary requirement, which means $D3$ and $D4$ must fire in ultrashort coincidence. In consequence most of the data collected at $D3$ and $D4$ must be discarded, as even for simultaneous emissions from the sources the mean value of τ_{34} is about $\sqrt{2}T$. This in turn precludes a test of local hidden variable theories by means of Bell's inequality at detectors $D3$ and $D4$, where a postselection throws away more than 31% of the data [21]. On the other hand, the test is possible at detectors $D1$ and $D2$ [Eq. (8)]. Again the requirement is for ultrashort coincidence at detectors $D3$ and $D4$ and not at $D1$ and $D2$, where there is no upper bound on the coincidence window $|t_1 - t_2|$. In fact, one would permit a wide coincidence window in order to collect all data at $D1$ and $D2$ that have been preselected by the ultrashort coincidence between $D3$ and $D4$. This constitutes a most discriminating test of Bell's inequality, because no postselection of the nonlocally correlated particles 1 and 2 is needed.

Let us now conclude with an attempt to understand how the polarization correlation of the particles on the left side can come about even when there are no polarizers on the right side and why only the mutual angles on each side—Eq. (4)—are relevant for the overall coincidences. The answer lies in the beam splitter. It superimposes the states of photons 3 and 4. The beams going to detectors $D3$ and $D4$ must therefore reflect the bosonic character of the particles and can only be occupied in the following two ways:

(a) *Both photons in the same beam.* The commutation rules demand that the two photons, e.g., from the beam going to $D3$ (assuming $r_1 = r_{11}, r_3 = r_4$, and $T_x = T_y = R_x = R_y = 1/2$) obey the following counterpart of Eq. (4):

$$P(\theta_1, \theta_2, 2 \times \theta_3) = \langle \Psi | \hat{E}_1^\dagger \hat{E}_2^\dagger \hat{E}_3^\dagger \hat{E}_3^\dagger \hat{E}_3 \hat{E}_3 \hat{E}_2 \hat{E}_1 | \Psi \rangle = \frac{1}{4} \cos^2(\theta_1 - \theta_3) \cos^2(\theta_2 - \theta_3) \quad (9)$$

given they have the same energy ($\omega_3 = \omega_4$). When there are no polarizers before $D3$ and $D4$, photons 1 and 2 going to the left would still have the polarization correlation, but individually they would be unpolarized. For both two-photon channels together we obtain $P(\theta_1, \theta_2, \infty, \infty) = [1 + \cos^2(\theta_1 - \theta_2)]/8$.

(b) *One photon in each beam: 1-1 channel.* Now the commutation rules require that the polarizations are

correlated as given by Eq. (4). This means we also get the correlation between the polarizations of beams 1 and 2 as given by Eq. (6): $P(\theta_1, \theta_2, \infty, \infty) = \sin^2(\theta_1 - \theta_2)/8$, i.e., with the maximum in the orthogonal direction with regard to the previous case.

Of course, the polarization correlation between photons 1 and 2 disappears if we do not detect them in coincidence detections on the right side, which one can see from the fact that the sum of $P(\theta_1, \theta_2, \infty, \infty)$ from all the channels is a constant.

To understand Eq. (4) let us compare it with the standard *left-right* Bell probabilities: $P(\theta_1, \theta_2) = \frac{1}{2} \cos^2(\theta_1 - \theta_2)$ and $P(\theta_3, \theta_4) = \frac{1}{2} \cos^2(\theta_3 - \theta_4)$. For the angles $\theta_1 = \theta_2$ and $\theta_3 = \theta_4$ we obtain [from Eq. (4)] $P(\theta_1, \theta_3, \theta_2, \theta_4) = 0$ no matter the values of $\theta_1 - \theta_3$ and $\theta_2 - \theta_4$. These three probabilities clearly cannot be satisfied simultaneously, and we can express this fact by saying that the fourth-order interference *erases* information on polarization correlation in the *1-1 channel*. An immediate consequence is that for $\theta_1 = \theta_2 = \theta_3 = \theta_4$ we can never register coincidences, and represents yet another possibility to formulate Bell's theorem without inequalities—the idea first developed by Greenberger, Horne, and Zeilinger [22]. The possibility is based on the well-known fact that the classical visibility of the fourth-order interference is not higher than 50%.

It is interesting that although for $\omega_3 \neq \omega_4$ we can track down the Bell *left-right* probabilities in the 2-photon beams exactly, as expected, in the *1-1 channel* this is even then not possible. The case of different energies can be handled with frequency filters (FF in Fig. 1). The result is then modulated with the beating period but in effect we have $P(\theta_1, \theta_2, \theta_3, \theta_4) = \frac{1}{2} \cos^2(\theta_1 - \theta_3) \cos^2(\theta_2 - \theta_4)$ if we assume two polarizers and two detectors in each two-photon beam (not shown in Fig. 1). So, a dropped polarizer in front of one of the four detectors immediately gives the standard *left-right* Bell probability for the other pair. The *1-1 channel*, on the other hand, responds to the special feature of the interference of the fourth order to “create” the polarization correlation even when unpolarized photons interfere—see Eq. (5).

Thus, while the second-order interference erases the *path* memory, the fourth-order interference erases the *polarization correlation memory*. It occurs in an analogous way in which the fourth-order interference erases the polarization memory of two polarized incident photons according to Eq. (16) of Ref. [17].

One of us (M.P.) is grateful to his host Professor K.-E. Hellwig, Institut für Theoretische Physik, TU Berlin and Prof. H. Paul, Humboldt-Universität zu Berlin for valuable discussions. He also acknowledges supports of the Alexander von Humboldt Foundation, the Technical University of Vienna, and the Ministry Science of Croatia.

Note added.—After one of us (M.P.) completed the first draft of the present paper containing the full elab-

oration in the plane waves (and not containing the ultrashort coincidence considerations introduced later on by the other author) we learned of Zukowski *et al.* [23] in which the *entanglement swapping* in the configuration space of down-converted polarized independent photons emerging from two nonlinear crystals was proposed together with the appropriate experimental conditions (ultracoincidence). We have also learned that such an *entanglement* was first recognized by Bennett *et al.* [24].

*Permanent address: Internet: pavicic@ati.ac.at

- [1] Z. Y. Ou, Phys. Rev. A **37**, 1607 (1988).
- [2] L. Mandel, Phys. Rev. A **28**, 929 (1983).
- [3] H. Paul, Rev. Mod. Phys. **57**, 209 (1986).
- [4] J. G. Rarity *et al.*, Phys. Rev. Lett. **65**, 1348 (1990).
- [5] Z. Y. Ou, X. Y. Zou, L. J. Wang, and L. Mandel, Phys. Rev. A **42**, 2957 (1990).
- [6] Z. Y. Ou, L. J. Wang, X. Y. Zou, and L. Mandel, Phys. Rev. A **41**, 566 (1990).
- [7] Z. Y. Ou and L. Mandel, Phys. Rev. Lett. **61**, 50 (1988).
- [8] X. Y. Ou, C. K. Hong, and L. Mandel, Opt. Commun. **67**, 159 (1988).
- [9] X. Y. Ou, C. K. Hong, and L. Mandel, Phys. Lett. A **122**, 11 (1987).
- [10] R. A. Campos, B. E. A. Saleh, and M. A. Teich, Phys. Rev. A **42**, 4127 (1990).
- [11] M. A. Horne, A. Shimony, and A. Zeilinger, Phys. Rev. Lett. **62**, 2209 (1989).
- [12] L. J. Wang, X. Y. Zou, and L. Mandel, Phys. Rev. Lett. **66**, 1111 (1991); See also X. Y. Zou *et al.*, Phys. Rev. Lett. **68**, 3667 (1992).
- [13] P. R. Holland and J. P. Vigiér, Phys. Rev. Lett. **67**, 402 (1991).
- [14] L. J. Wang, X. Y. Zou, and L. Mandel, Phys. Rev. Lett. **67**, 403 (1991).
- [15] M. P. Silverman, Am. J. Phys. **61**, 514 (1993).
- [16] J. S. Bell, Phys. Rep. **137**, 7 (1986).
- [17] M. Pavičić, Phys. Rev. A **50**, 3486 (1994).
- [18] A. Aspect, J. Dalibard, and G. Roger, Phys. Rev. Lett. **49**, 1804 (1982).
- [19] For the given operators cf. Refs. [1,6–8]. Note that positive imaginary terms assure the preservation of boson commutation relations for the annihilation operators at the output of the beam splitter and that Eqs. (2), (5a), etc. of Ref. [7] and Eqs. (56), (57), etc. of Ref. [1] should be corrected accordingly. We would like to mention here that Professor H. Paul noticed the latter fact as well (private communication).
- [20] B. Yurke and D. Stoler, Phys. Rev. Lett. **68**, 1251 (1992).
- [21] A. Garg and N. D. Mermin, Phys. Rev. D **35**, 3831 (1987).
- [22] D. M. Greenberger, M. Horne, and A. Zeilinger, in *Bell's Theorem, Quantum Theory, and Conceptions of the Universe*, edited by M. Kafatos (Kluwer Academic, Dordrecht, The Netherlands, 1989), p. 73.
- [23] M. Zukowski, A. Zeilinger, M. A. Horne, and A. K. Ekert, Phys. Rev. Lett. **71**, 4287 (1993).
- [24] C. H. Bennett, C. Brassard, C. Crépeau, R. Jozsa, A. Peres, and W. K. Wootters, Phys. Rev. Lett. **70**, 1895 (1993).