

Orthomodular Lattices and a Quantum Algebra

Norman D. Megill^{1,2} and Mladen Pavičić^{3,4}

Received February 4, 2001

We show that one can formulate an algebra with lattice ordering so as to contain one quantum and five classical operations as opposed to the standard formulation of the Hilbert space subspace algebra. The standard orthomodular lattice is embeddable into the algebra. To obtain this result we devised algorithms and computer programs for obtaining expressions of all quantum and classical operations within an orthomodular lattice in terms of each other, many of which are presented in the paper. For quantum disjunction and conjunction we prove their associativity in an orthomodular lattice for any triple in which one of the elements commutes with the other two and their distributivity for any triple in which a particular element commutes with the other two. We also prove that the distributivity of symmetric identity holds in Hilbert space, although whether or not it holds in all orthomodular lattices remains an open problem, as it does not fail in any of over 50 million Greechie diagrams we tested.

1. INTRODUCTION

Closed subspaces of Hilbert space form an algebra called a Hilbert lattice. A Hilbert lattice is a kind of orthomodular lattice, which we, in the next section, introduce starting with an ortholattice, which is a still simpler structure. In any Hilbert lattice the operation *meet*, $a \cap b$, corresponds to set intersection $\mathcal{H}_a \cap \mathcal{H}_b$ of subspaces \mathcal{H}_a and \mathcal{H}_b of Hilbert space \mathcal{H} ; the ordering relation $a \leq b$ corresponds to $\mathcal{H}_a \subseteq \mathcal{H}_b$; the operation *join*, $a \cup b$, corresponds to the smallest closed subspace of \mathcal{H} containing $\mathcal{H}_a \cup \mathcal{H}_b$; and a' corresponds to \mathcal{H}'_a , the set of vectors orthogonal to all vectors in \mathcal{H}_a . Within Hilbert space there is also an operation that has no parallel in the Hilbert lattice: the sum of two subspaces $\mathcal{H}_a + \mathcal{H}_b$, which is defined as the set of sums of vectors from \mathcal{H}_a and \mathcal{H}_b . We

¹ Boston Information Group, 30 Church St., Belmont MA 02478, U.S.A.

² E-mail: nm@alum.mit.edu; Web page: <http://www.metamath.org>

³ University of Zagreb, Gradjevinski Fakultet, Kačićeva 26, HR-10000 Zagreb, Croatia.

⁴ E-mail: mpavicic@faust.irb.hr; Web page: <http://m3k.grad.hr/pavicic>

also have, $\mathcal{H}_a + \mathcal{H}'_a = \mathcal{H}$. One can define all the lattice operations on Hilbert space itself, following the above definitions ($\mathcal{H}_a \cap \mathcal{H}_b = \mathcal{H}_a \cap \mathcal{H}_b$, etc.). Thus we have $\mathcal{H}_a \cup \mathcal{H}_b = \overline{\mathcal{H}_a + \mathcal{H}_b} = (\mathcal{H}_a + \mathcal{H}_b)^{\perp\perp} = (\mathcal{H}'_a \cap \mathcal{H}'_b)'$, (Isham, 1995, p. 175) where $\overline{\mathcal{H}_c}$ is a closure of \mathcal{H}_c , and therefore $\mathcal{H}_a + \mathcal{H}_b \subseteq \mathcal{H}_a \cup \mathcal{H}_b$. When \mathcal{H} is finite dimensional or when the closed subspaces \mathcal{H}_a and \mathcal{H}_b are orthogonal to each other, then $\mathcal{H}_a + \mathcal{H}_b = \mathcal{H}_a \cup \mathcal{H}_b$ (Halmos, 1995, pp. 21–29; Kalmbach, 1983, pp. 66–67; Mittelstaedt, 1978, pp. 8–16).

In the past, scientists, starting with Birkhoff and von Neumann, wanted to find parallels with a possible logic lying underneath the orthomodular lattice and operations defined on such a logic. A possible candidate for the logic was formulated (Chiara, 1986; Dishkant, 1974; Kalmbach, 1974; Kalmbach, 1983; Mittelstaedt, 1978). However, it has recently been shown (Pavičić and Megill, 1998a) that the logic can have at least two models: Hilbert space and another model that is not orthomodular—so there is no *proper* quantum logic.⁵ One can still consider operations within the model itself: the orthomodular lattice. The problem of finding quantum operations that would reduce to classical ones for compatible observables has been attacked many times in the past. In particular, it has been shown that one can start with unique *classical* conjunction, disjunction, and implication and using them define five *quantum* conjunctions, disjunctions, and implications [which collapse into former classical ones for commuting (compatible, commensurable) observables]. In this paper we show that one can start with unique quantum operations and arrive at five classical ones. Thus it turns out that the usual way of defining orthomodular lattice by means of unique classical conjunction and disjunction is a consequence of a direct translation of meet and join from Hilbert space. We also express all possible quantum and classical operations by each other, even a chosen classical or quantum one by means of all other quantum and classical ones in single equations. We do so with the help of a computer program that reduces two-variable expressions to each other.

In section 5 we prove that in an orthomodular lattice the associativity of both quantum disjunctions and conjunctions holds for any triple of lattice elements as soon as one of them commutes with the other two.

In the end, we partially solve an open problem from Megill and Pavičić (2000) by proving that the “distributive law” for a quantum identity holds in the Godowski lattices and therefore in Hilbert space. It remains an open problem whether or not the law holds in all orthomodular lattices.

⁵ Consequently, the papers that are now appearing and claim—as, e.g., Dalla Chiara and Giuntini (2001)—that quantum logic, defined as a genuine logical system, characterizes orthomodular lattices are simply incorrect. All previous such papers and books are outdated by the result.

2. QUANTUM AND CLASSICAL LATTICE OPERATIONS

One usually defines an ortholattice in the following way.

Definition 2.1. An ortholattice (OL) is an algebra $\langle \mathcal{L}_0, ', \cup \rangle$ such that the following conditions are satisfied for any $a, b, c \in \mathcal{L}_0$:

$$L1 \quad a = a'' \tag{2.1}$$

$$L2 \quad a \leq a \cup b \quad \& \quad b \leq a \cup b \quad \& \quad b \leq a \cup a' \tag{2.2}$$

$$L3 \quad a \leq b \quad \& \quad b \leq a \Rightarrow a = b; \quad a = b \Rightarrow a \leq b \tag{2.3}$$

$$L4 \quad a \leq b \Rightarrow b' \leq a' \tag{2.4}$$

$$L5 \quad a \leq b \quad \& \quad b \leq c \Rightarrow a \leq c \tag{2.5}$$

$$L6 \quad a \leq c \quad \& \quad b \leq c \Rightarrow a \cup b \leq c \tag{2.6}$$

where

$$a \leq b \stackrel{\text{def}}{=} a \cup b = b, \quad 1 \stackrel{\text{def}}{=} a \cup a', \quad 0 \stackrel{\text{def}}{=} a \cap a'. \tag{2.7}$$

Then we can define six operations of implication:

Definition 2.2. $a \rightarrow_0 b \stackrel{\text{def}}{=} a' \cup b, a \rightarrow_1 b \stackrel{\text{def}}{=} a' \cup (a \cap b), a \rightarrow_2 b \stackrel{\text{def}}{=} b' \rightarrow_1 a', a \rightarrow_3 b \stackrel{\text{def}}{=} ((a' \cap b) \cup (a' \cap b')) \cup (a \cap (a' \cup b)), a \rightarrow_4 b \stackrel{\text{def}}{=} a' \rightarrow_3 b', a \rightarrow_5 b \stackrel{\text{def}}{=} ((a \cap b) \cup (a' \cap b)) \cup (a' \cup b')$, where \rightarrow_0 is called *classical implication* and $\rightarrow_i, i = 1, \dots, 5$ quantum implication.

Quantum implications reduce to the classical one whenever a and b commute.

Definition 2.3. We say that a and b commute and write aCb when any and therefore all of the following equations hold: (Holland, 1995; Mittelstaedt, 1978; Zeman, 1979) $(a \cap b) \cup (a \cap b') \cup (a' \cap b) \cup (a' \cap b') = 1, a \cap (a' \cup b) \leq b, a = (a \cap b) \cup (a \cap b')$.

We can also define:

Definition 2.4.

$$a \cup_i b \stackrel{\text{def}}{=} a' \rightarrow_i b, \quad a \cap_i b \stackrel{\text{def}}{=} (a \rightarrow_i b)', \quad i = 0, \dots, 5; \tag{2.8}$$

$$a \equiv_i b \stackrel{\text{def}}{=} (a \rightarrow_i b) \cap (b \rightarrow_0 a), \quad i = 0, \dots, 5, \tag{2.9}$$

where $a \cup_0 b = a \cup b, a \cap_0 b = a \cap b$ and $a \equiv_0 b$ are classical disjunction, conjunction, and identity, respectively, while $a \cup_i b, a \cap_i b$, and $a \equiv_i b, i = 1, \dots, 5$,

are quantum ones, respectively. The latter obviously reduce to the former when a and b commute.

For the above operations the following theorems hold. In them, we can also pick any one of the conditions in Theorem 2.5 as our definition of an orthomodular lattice and in Theorem 2.6 as our definition of a distributive lattice (Boolean algebra).

Theorem 2.5. *An ortholattice in which any one of the following conditions holds is an orthomodular lattice and vice versa (Pavičić, 1987, 1989, 1993, 1998; Pavičić and Megill, 1998b).*

$$a \rightarrow_i b = 1 \Leftrightarrow a \leq b, \quad i = 1, \dots, 5, \quad (2.10)$$

$$a \cup_i b = 1 \Leftrightarrow a' \perp b', \quad i = 1, \dots, 5, \quad (2.11)$$

$$a \cap_i b = 0 \Leftrightarrow a \perp b, \quad i = 1, \dots, 5, \quad (2.12)$$

$$a \equiv_i b = 1 \Leftrightarrow a = b, \quad i = 1, \dots, 5, \quad (2.13)$$

$$a \perp b \ \& \ a \cup b = 1 \Rightarrow a' \perp b', \quad (2.14)$$

where $a \perp b \stackrel{\text{def}}{=} a \leq b'$

Theorem 2.6. *An ortholattice in which any one of the following conditions holds is a distributive lattice and vice versa (Pavičić, 1987, 1989, 1993, 1998; Pavičić and Megill, 1998b).*

$$a \rightarrow_0 b = 1 \Leftrightarrow a \leq b, \quad (2.15)$$

$$a \cup b = a \cup_0 b = 1 \Leftrightarrow a' \perp b', \quad (2.16)$$

$$a \cap b = a \cap_0 b = 0 \Leftrightarrow a \perp b, \quad (2.17)$$

$$a \equiv_0 b = 1 \Leftrightarrow a = b. \quad (2.18)$$

Actually, in any orthomodular lattice all expressions with two variables are reducible to 1 of the 96 Beran canonical forms. (Beran, 1985, Table 1, p. 82) The reader can easily reduce any two-variable expression with the help of our program `beran`, which we describe in section 8. All 96 forms can be also viewed inside the source code of `beran.c`. In general we can divide them into 16 *classical* and 80 *quantum* ones. Classical expressions are, classical implication and its negation (disjunction and conjunction) – Beran expressions 2–5 and 92–95; classical identity and its negation – expressions 9 and 88; variables a , b and their negations – expressions 22, 39, 58, and 75; and “0” and “1” – expressions 1 and 96, respectively. Quantum expressions are all the other expressions, that reduce to classical ones whenever the variables commute: quantum implications and their negations (quantum disjunctions and conjunctions) – 12–15, 18–21, 28–31, 34–37,

44–47, 50–53, 60–63, 66–69, 76–79, and 82–85; quantum identities ($a \equiv_1 b = a' \equiv_3 b'$, $a \equiv_2 b = a' \equiv_4 b'$, $a \equiv_5 b$) and their negations – 24, 25, 40, 41, 56, 57, 72, 73, 8, and 89; “quantum variables” (which reduce to “classical a, b ”) and their negations – a : 6, 38, 54, 70, 86, b : 7, 23, 55, 71, 87, $-a$: 11, 27, 43, 59, 91, and $-b$: 10, 26, 42, 74, 90; and “quantum 0,1” – “0”: 17, 33, 49, 65, 81 and “1”: 16, 32, 48, 64, 80. For some of these quantum expressions we give the following definitions and theorems.

Definition 2.7. Quantum unities and zeros in an OML are,

$$1_{1(a,b)} \stackrel{\text{def}}{=} a' \cup (a \cap b') \cup (a \cap b) \tag{2.19}$$

$$1_{2(a,b)} \stackrel{\text{def}}{=} b \cup (a \cap b') \cup (a' \cap b') \tag{2.20}$$

$$1_{3(a,b)} \stackrel{\text{def}}{=} a \cup (a' \cap b) \cup (a' \cap b') \tag{2.21}$$

$$1_{4(a,b)} \stackrel{\text{def}}{=} b' \cup (a' \cap b) \cup (a \cap b) \tag{2.22}$$

$$1_{5(a,b)} \stackrel{\text{def}}{=} (a \cap b) \cup (a \cap b') \cup (a' \cap b) \cup (a' \cap b') \tag{2.23}$$

$$0_{1(a,b)} \stackrel{\text{def}}{=} a \cap (a' \cup b) \cap (a' \cup b') \tag{2.24}$$

$$0_{2(a,b)} \stackrel{\text{def}}{=} b' \cap (a' \cup b) \cap (a \cup b) \tag{2.25}$$

$$0_{3(a,b)} \stackrel{\text{def}}{=} a' \cap (a \cup b') \cap (a \cup b) \tag{2.26}$$

$$0_{4(a,b)} \stackrel{\text{def}}{=} b \cap (a \cup b') \cap (a' \cup b') \tag{2.27}$$

$$0_{5(a,b)} \stackrel{\text{def}}{=} (a \cup b) \cap (a \cup b') \cap (a' \cup b) \cap (a' \cup b') \tag{2.28}$$

Some consequences of these definitions are straightforward:

Lemma 2.8. *Two variables commute iff any of the 80 two-variable quantum expressions is equal to its classical counterpart. Two variables also commute iff any two of the five different forms of each quantum expression are equal to each other.*

For example, $1_{i(a,b)} = 1$ or $0_{i(a,b)} = 0$, $i = 1, \dots, 5$ is equivalent to aCb . In particular, $1_{5(a,b)} = 1$ is the first expression from Definition 2.3. Also, for example, $a_{i(a,b)} = a$, $i = 1, \dots, 5$, where $a_{i(a,b)}$ are given by Beran expressions: 6, 38, 54, 70, 86, respectively, are equivalent to aCb . In particular, $a_{1(a,b)} = (a \cap b) \cup (a \cap b') = a$ is the third expression from Definition 2.3. Examples for the second claim of the theorem are that $a \rightarrow_i b = a \rightarrow_j b$, $a \cup_i b = a \cup_j b$, and $a \cap_i b = a \cap_j b$, $i \neq j$, $i, j = 1, \dots, 5$ are equivalent to aCb (Pavičić, 1993, p. 1487). The same holds for $1_{i(a,b)} = 1_{j(a,b)}$, $a_{i(a,b)} = a_{j(a,b)}$, $a \equiv_i b = a \equiv_j b$, $i \neq j$, $i, j = 1, \dots, 5$, etc.

Theorem 2.9. *An ortholattice in which any one of the following conditions holds is an orthomodular lattice and vice versa.*

$$a \rightarrow_i b = 1_{j(a,b)} \Leftrightarrow a \leq b, \quad i, j = 1, \dots, 5; \quad i \neq j; \quad (2.29)$$

$$a \equiv_i b = 1_{j(a,b)} \Leftrightarrow a = b, \quad i, j = 1, \dots, 5; \quad i \neq j. \quad (2.30)$$

Proof. We will exemplify the proofs by proving the case $i = 1, j = 5$. Other cases the reader can prove analogously. We first use Eq. (2.13) to write the premise as $(a \rightarrow_1 b) \equiv_5 1_5 = 1$ (\equiv_5 should be used for all cases—in it the subscript 5 is not j) and then we find the canonical expression of

$$(a \rightarrow_1 b) \equiv_5 1_{5(a,b)} = (a \rightarrow_1 b) \equiv_5 (a \cap b) \cup (a \cap b') \cup (a' \cap b) \cup (a' \cap b')$$

by typing (see section 8 for details on our program `beran`)

$$\text{beran “((aIb) = (((a^b)v(a^b))v((a^b)v(a^b))))”$$

The program responds with

$$30 ((-avb)^(av(-a^b))v(-a^b))$$

which is nothing but $a \rightarrow_3 b$. Using Eq. (2.10) we get the desired conclusion. \square

3. RELATIONS BETWEEN OPERATIONS

In this section we show how one can connect the operations defined in section 2 with each other in an orthomodular lattice defined in a standard way given by Definition 2.1. In counting the cases for commuting operations later, we disregard the order of a and b .

In Pavičić and Megill (1998b) we have shown how one can express classical disjunction by quantum and classical implications within a single equation. (That equation was one of the four smallest ones. Another one is presented later.)

Lemma 3.1. (i) *The equation*

$$a \cup b = (((b \rightarrow_i a) \rightarrow_i (a \rightarrow_i b)) \rightarrow_i b) \rightarrow_i a \rightarrow_i a \quad (3.1)$$

is true in all orthomodular lattices for $i = 1, \dots, 5$ and in all distributive lattices for $i = 0, \dots, 5$;

(ii) *an ortholattice in which Eq. 3.1 holds is an orthomodular lattice for $i = 1, \dots, 5$ and a distributive lattice for $i = 0$.*

This equation does not contain negations and if we wanted to define an algebra by means of so merged implications and without using negation we should at least introduce 0. Alternatively one can use the negation and define 0. In this paper we adopt the latter approach. We do not give proofs of the lemmas in this section because all expressions can be trivially checked with the help of the computer

program *beran* written by one of us (N. D. M.), which the reader can download from our web sites.

Lemma 3.2. *There is only one “smallest” (lowest number of occurrence of variables, 5, and negations, 2) expression of classical disjunction by means of quantum implications:*

$$a \cup b = (((a' \rightarrow_i b') \rightarrow_i b) \rightarrow_i a) \rightarrow_i a; \quad i = 1, \dots, 5, \quad (3.2)$$

and seven smallest (5 variables, 4 negations) expressions of classical conjunction by means of quantum implications, one of which is,

$$a \cap b = (a \rightarrow_i ((a \rightarrow_i b) \rightarrow_i (b' \rightarrow_i a')'))'; \quad i = 1, \dots, 5. \quad (3.3)$$

There are two smallest (5 variables, 3 negations) expressions of classical disjunction by means of quantum disjunctions, one of which is,

$$a \cup b = ((a \cup_i b') \cup_i (b' \cup_i a))' \cup_i a; \quad i = 1, \dots, 5, \quad (3.4)$$

and two (5,5) by means of quantum conjunctions, one of which is,

$$a \cup b = (((a' \cap_i b) \cap_i (b \cap_i a'))' \cap_i a')'; \quad i = 1, \dots, 5, \quad (3.5)$$

An equal number of smallest expressions of classical conjunction by means of quantum disjunctions and conjunctions we get by using $a \cap b = (a' \cup b')'$ and $a \cap_i b = (a' \cup_i b')'$ (of course with reversed smallest number of negations).

Any of these equations when added to an ortholattice makes it orthomodular.

Lemma 3.3. *Here are samples of the smallest expressions (with their numbers being given in curly brackets) of classical conjunction and disjunction by means of both, classical ($i = 0$) and quantum ($i = 1, \dots, 5$) implications, disjunctions, and conjunctions in single equations in any orthomodular lattice:*

$$a \cup b = ((b \rightarrow_i a) \rightarrow_i (((a \rightarrow_i b') \rightarrow_i b') \rightarrow_i a)) \quad \{1\} \quad (3.6)$$

$$a \cup b = (b \cup_i (a \cup_i ((a \cup_i b) \cup_i (b' \cup_i a)))) \quad \{16\} \quad (3.7)$$

$$a \cup b = ((a' \cap_i b') \cap_i (a' \cap_i (b \cap_i (b \cap_i a'))))' \quad \{8\} \quad (3.8)$$

$$a \cap b = (a \rightarrow_i ((a \rightarrow_i ((a \rightarrow_i b) \rightarrow_i b')) \rightarrow_i a'))' \quad \{23\} \quad (3.9)$$

$$a \cap b = ((b' \cup_i (a \cup_i (a \cup_i b')) \cup_i (b' \cup_i a))' \quad \{8\} \quad (3.10)$$

$$a \cap b = (b \cap_i (a \cap_i ((a \cap_i b) \cap_i (b' \cap_i a)))) \quad \{16\}, \quad (3.11)$$

where $i = 0, \dots, 5$. Any of these equations for $i = 1, \dots, 5$ and Eqs. (3.6), (3.8), (3.9), and (3.10) for $i = 0$ when added to an ortholattice makes it orthomodular (fails in O6). For $i = 0$, there are no such smallest samples of the type given by Eqs. (3.7) and (3.11) and there are 18 samples that pass O6 of Eq. (3.9) type, 4 of

(3.8) type, and 4 of (3.10) type. Samples of the latter ones are,

$$a \cap b = (a \rightarrow_i (b \rightarrow_i ((b \rightarrow_i a) \rightarrow_i (b' \rightarrow_i a'))))' \tag{3.12}$$

$$a \cup b = (b' \cap_i (a' \cap_i ((a' \cap_i b) \cap_i (b \cap_i a'))))' \tag{3.13}$$

$$a \cap b = (b' \cup_i (a' \cup_i ((a' \cup_i b) \cup_i (b \cup_i a'))))', \tag{3.14}$$

respectively.

Lemma 3.4. *The shortest expressions of some operations defined here by each other are,*

$$\begin{aligned} a \cup b &= a \cup_0 b = b \cup_1 (b \cup_1 a') = a \cup_2 (b' \cup_2 a') = b \cup_3 (b \cup_3 a) \\ &= a \cup_4 (b \cup_4 a) = b \cup_5 (b \cup_5 a') \end{aligned} \tag{3.15}$$

$$a \cup_1 b = b \cup_2 a = (a \cup_3 b) \cup_3 b = b \cup_4 (b \cup_4 a) = a \cup_5 (b \cup_5 a) \tag{3.16}$$

$$a \cup_2 b = b \cup_1 a = (b \cup_3 a) \cup_3 a = a \cup_4 (a \cup_4 b) = b \cup_5 (b \cup_5 a) \tag{3.17}$$

$$\begin{aligned} a \cup_3 b &= b \cup_4 a = (a \cup_1 b') \cup_1 (b \cup_1 a) = (a \cup_2 b) \cup_2 (b' \cup_2 a') \\ &= (a \cup_5 b) \cup_5 (a \cup_5 (b \cup_5 a'))' \end{aligned} \tag{3.18}$$

$$\begin{aligned} a \cup_4 b &= b \cup_3 a = (b \cup_1 a') \cup_1 (a \cup_1 b) = (b \cup_2 a) \cup_2 (a' \cup_2 b') \\ &= (b \cup_5 a) \cup_5 (b \cup_5 (a \cup_5 b'))' \end{aligned} \tag{3.19}$$

$$\begin{aligned} a \cup_5 b &= b \cup_5 a = ((a \cup_1 b)' \cup_1 (a' \cup_1 (b \cup_1 a)))' \\ &= ((b \cup_2 a)' \cup_2 ((a \cup_2 b) \cup_2 a'))' \\ &= ((b \cup_3 a)' \cup_3 ((b \cup_3 a) \cup_3 a'))' \\ &= ((b \cup_4 a)' \cup_4 (b \cup_4 (b \cup_4 a)))' \end{aligned} \tag{3.20}$$

$$a \equiv_0 b = (a' \equiv_5 b') = (b \cup_i a)' \cup_i (b' \cup_i a)'; \quad i = 1, \dots, 5 \tag{3.21}$$

$$\begin{aligned} a \equiv_1 b &= a' \equiv_3 b' = (a \cup_{1,3} b)' \cup_{1,3} (b' \cup_{1,3} a) \\ &= (a' \cup_{2,4} b')' \cup_{2,4} (b \cup_{2,4} a) \\ &= (a \cup_5 (a \cup_5 b))' \cup_5 (a' \cup_5 (b \cup_5 a'))' \end{aligned} \tag{3.22}$$

$$\begin{aligned} a \equiv_2 b &= a' \equiv_4 b' = (b' \cup_{1,3} a)' \cup_{1,3} (a \cup_{1,3} (b \cup_{1,3} a)) \\ &= ((a' \cup_2 b)' \cup_2 (a \cup_2 (b \cup_2 a)))' \\ &= (a \cup_4 b)' \cup_4 (a' \cup_4 (b' \cup_4 a))' \\ &= ((b \cup_5 (b \cup_5 a))' \cup_5 (a \cup_5 (b \cup_5 a')))' \end{aligned} \tag{3.23}$$

Dual expressions on both sides of equations we get by using $a \cap_i b = (a' \cup_i b)'$.

Lemma 3.5. *Samples of expressions of particular quantum disjunctions by means of all five of them together in single equations are*

$$a \cup_1 b = a \cup_i (b' \cup_i (b \cup_i a)') \tag{3.24}$$

$$a \cup_2 b = b \cup_i (a' \cup_i (a \cup_i b)') \tag{3.25}$$

$$a \cup_3 b = ((a' \cup_i (b \cup_i a))' \cup_i ((b \cup_i (a \cup_i b)) \cup_i (b' \cup_i a)'))' \tag{3.26}$$

$$a \cup_4 b = ((b' \cup_i (a \cup_i b))' \cup_i ((a \cup_i (b \cup_i a)) \cup_i (a' \cup_i b)'))' \tag{3.27}$$

$$a \cup_5 b = ((b \cup_i a)' \cup_i (b' \cup_i ((a \cup_i b) \cup_i (b \cup_i a)')))' \tag{3.28}$$

where $i = 1, \dots, 5$. Dual expressions ($a \cup b$ by means of $a \cap_i b$, and $a \cap b$ by means of $a \cup_i b$ and $a \cap_i b$) we get by using $a \cap b = (a' \cup b)'$ and $a \cap_i b = (a' \cup_i b)'$.

4. QUANTUM ALGEBRA

In Lemma 3.2, Eq. (3.4), we have shown how one can express the classical disjunction by means of quantum ones in a single equation. So, we can substitute this expression for the disjunctions in conditions that define an orthomodular lattice (Definition 2.1) and obtain five formally identical ways to write those conditions using five quantum disjunctions. But we can do even more and define an algebra with a lattice ordering as follows.

Definition 4.1. A quantum algebra QA is an algebra $\langle \mathcal{A}_O, ', \cup \rangle$ such that the following conditions are satisfied for any $a, b, c \in \mathcal{A}_O$:

$$A1 \quad a = a'' \quad \& \quad b \leq 1 \tag{4.1}$$

$$A2 \quad a \leq ((a \cup b') \cup (b' \cup a))' \cup a \quad \& \quad b \leq ((a \cup b') \cup (b' \cup a))' \cup a \tag{4.2}$$

$$A3 \quad a \leq b \quad \& \quad b \leq a \Rightarrow a = b; \quad a = b \Rightarrow a \leq b \tag{4.3}$$

$$A4 \quad a \leq b \Rightarrow b' \leq a' \tag{4.4}$$

$$A5 \quad a \leq b \quad \& \quad b \leq c \Rightarrow a \leq c \tag{4.5}$$

$$A6 \quad a \leq c \quad \& \quad b \leq c \Rightarrow ((a \cup b') \cup (b' \cup a))' \cup a \leq c \tag{4.6}$$

$$A7 \quad a \perp b \quad \& \quad ((a \cup b') \cup (b' \cup a))' \cup a = 1 \Rightarrow a' \perp b', \tag{4.7}$$

where

$$a \leq b \stackrel{\text{def}}{=} ((a \cup b') \cup (b' \cup a))' \cup a = b \tag{4.8}$$

$$1 \stackrel{\text{def}}{=} ((a \cup a) \cup (a \cup a))' \cup a \quad \& \quad 0 \stackrel{\text{def}}{=} (((a \cup a) \cup (a \cup a))' \cup a)'. \tag{4.9}$$

Substitution Rule: Any valid condition or equation one can obtain in the standard formulation of OML containing only variables, \cup_i (satisfied for all $i = 1, \dots, 5$), and negations written in QA with \cup substituted for \cup_i is a valid condition or equation in QA.

We can easily check that the above ordering is a proper ordering and that for $a, b, c \in \mathcal{A}_0$ lower upper and greater lower bounds exist – they are given by $((a \cup b') \cup (b' \cup a))' \cup a$ and $((a' \cup b) \cup (b \cup a'))' \cup a'$, respectively. Obviously we can introduce the following definition $x \cup y \stackrel{\text{def}}{=} ((a \cup b') \cup (b' \cup a))' \cup a$ and obtain the standard definition of OML as given in section 2. This enables us to formulate the above *Substitution Rule*, which actually introduces an infinite number of conditions. Whether or not they can be replaced with a finite set of individual conditions is an open problem. Along this rule, A7 becomes Eq. (2.14). Equation (2.14) is equivalent to Eq. (2.13), which for $j = 5$ reads (Pavičić and Megill, 1999): $a \equiv_5 b = (a \cap b) \cup (a' \cap b') = 1 \Leftrightarrow a = b$. Since from Eq. (3.23) we have $a \equiv_5 b = ((b \cup_i a') \cup_i (b' \cup a))'$, $i = 1, \dots, 5$, we get A8. Similarly, we get A9, etc. Of course, we can never arrive at $a \leq a \cup b$, $a \cup (a \cap b) = a$, $a \cup (a' \cap (a \cup b)) = a \cup b$, or many other equations we are used to in OML. For example, if we had had $a \cup (a \cap b) = a$, that would have reduced Eq. (4.12) to Eq. (2.17) and therefore turn QA into a Boolean algebra.

Lemma 4.2.

$$\text{A8} \quad (b \cup a') \cap (b' \cup a) = 1 \Leftrightarrow a = b, \quad (4.10)$$

$$\text{A9} \quad a \cup a' = 1, \quad (4.11)$$

$$\text{A10} \quad a \cap (b \cup (b \cap a)) = 0 \Leftrightarrow a \perp b, \quad (4.12)$$

$$\text{A11} \quad (((b \cup a) \cap a') \cup b) \cap a = a \cap (b \cup (b \cap a)), \quad (4.13)$$

$$\text{A12} \quad a \cup b = 1 \Leftrightarrow a' \perp b', \quad (4.14)$$

where $a \cap b \stackrel{\text{def}}{=} (a' \cup b')$.

On the other hand, Lemma 3.4 indicates that there might be different ways of expressing classical disjunctions by means of quantum ones. And indeed, $a \cup b = (a \cup_5 b) \cup_5 (b' \cup_5 a)'$ does not match any other \cup_i – meaning $a \cup b \neq (a \cup_i b) \cup_i (b' \cup_i a)'$, $i = 1, 2, 3, 4$. The same is true with Eq. (3.15) for \cup_3 and \cup_4 , as well as with $a \cup b = ((a' \cup_1 b') \cup_1 b') \cup_1 ((a \cup_1 b') \cup_1 a')$ and $a \cup b = ((a' \cup_2 b') \cup_2 a') \cup_2 (b' \cup_2 (b \cup_2 a'))'$. Thus we arrive at the following theorem.

Theorem 4.3. *In QA one can express classical disjunction in the following five nonequivalent ways:*

$$a \cup_{cl1} b = ((a' \cup b') \cup b') \cup ((a \cup b') \cup a')' \quad (4.15)$$

$$a \cup_{cl2} b = ((a' \cup b') \cup a') \cup (b' \cup (b \cup a'))' \quad (4.16)$$

$$a \cup_{cl3} b = b \cup (b \cup a) \tag{4.17}$$

$$a \cup_{cl4} b = a \cup (b \cup a) \tag{4.18}$$

$$a \cup_{cl5} b = (a \cup b) \cup (b' \cup a)'. \tag{4.19}$$

Of course, there are many other such nonequivalent 5-tuples. Altogether, there are $(5^5)^5$ 5-tuples.

In conclusion, by using the parallels with the standard orthomodular lattice theory, in QA we can derive all the equations that hold in the lattice theory in terms of $\cup_i, i = 1, \dots, 5$, and negation, even those that cannot be obtained by the method presented in section 3 – for example, $a \cup_i (b \cap_i a) = a \cup_i (b' \cap_i a)$ or $a \cup_i (b \cup_i (a' \cap_i (a \cup_i b))) = a \cup_i b$ where neither side of these equations are equal to particular Beran expressions for all $i = 1, \dots, 5$, while the equations themselves do hold for all $i = 1, \dots, 5$. On the other hand, by using $a \cup b \stackrel{\text{def}}{=} ((a \cup b') \cup (b' \cup a))' \cup a$ and A1–A7 from Definition 4.1 we can embed the standard orthomodular lattice theory in QA.

5. CONDITIONAL ASSOCIATIVITY OF QUANTUM OPERATIONS

Quantum disjunctions and conjunctions are not associative. However, a conditional associativity, similar to Foulis–Holland (F–H) distributivity, does hold in any orthomodular lattice as proved in the theorem later. D’Hooghe and Pykacz (2000, p. 648) proved the theorem for $i = 1, 2$, and 5 , and conjectured it for $i = 3$ and 4 . Here we confirm their conjecture by giving the proofs for $i = 3$ and 4 . By doing so we prove that the conditional associativity holds for the unified quantum disjunction and conjunction (\cup and \cap) from the previous section. For this purpose, we may take aCb to be $a \cup (a' \cap b) = b \cup a$, noting that in any OML $a \cup_i (a' \cap_i b) = b \cup_i a$ is equivalent to aCb for $i = 1, \dots, 5$.

Theorem 5.1. *In any orthomodular lattice any triple $\{a, b, c\}$ in which one of the elements commutes with the other two is associative with respect to \cup_i and $\cap_i, i = 1, \dots, 5$:*

$$aCb \ \& \ aCc \Rightarrow (a \cup_i b) \cup_i c = a \cup_i (b \cup_i c), \quad i = 1, \dots, 5 \tag{5.1}$$

$$aCb \ \& \ bCc \Rightarrow (a \cup_i b) \cup_i c = a \cup_i (b \cup_i c), \quad i = 1, \dots, 5 \tag{5.2}$$

$$aCc \ \& \ bCc \Rightarrow (a \cup_i b) \cup_i c = a \cup_i (b \cup_i c), \quad i = 1, \dots, 5 \tag{5.3}$$

$$aCb \ \& \ aCc \Rightarrow (a \cap_i b) \cap_i c = a \cap_i (b \cap_i c), \quad i = 1, \dots, 5 \tag{5.4}$$

$$aCb \ \& \ bCc \Rightarrow (a \cap_i b) \cap_i c = a \cap_i (b \cap_i c), \quad i = 1, \dots, 5 \tag{5.5}$$

$$aCc \ \& \ bCc \Rightarrow (a \cap_i b) \cap_i c = a \cap_i (b \cap_i c), \quad i = 1, \dots, 5 \tag{5.6}$$

Proof. Since D'Hooghe and Pykacz (2000, p. 648) proved the cases $i = 1, 2, 5$ we only give sketchy proofs for these cases for the sake of completeness.

For $i = 1$, Eq. (5.1), given the premise aCb and the F–H theorem [$aCb \& aCc \Rightarrow (a \cup b) \cap c = (a \cap c) \cup (b \cup c)$, etc] we have (since $a'Ca$): $a \cup_1 b = a \cup (a' \cap b) = (a \cup a') \cap (a \cup b) = a \cup b$. Thus, the conclusion from Eq. (5.1) reads

$$(a \cup_1 b) \cup_1 c = a \cup b \cup c = a \cup_1 (b \cup_1 c).$$

Equations (5.2) and (5.3) follow analogously. Since $a \cup_2 b = b \cup_1 a$ and $a \cap_{1,2} b = (a' \cup_{1,2} b)'$, we have proved the theorem for $i = 1, 2$.

For $i = 5$, (again we have $aCb \Rightarrow a \cup_5 b = a \cup b$, etc.) both sides of the conclusion of Eqs. (5.1), (5.2), and (5.3) reduce to $a \cup (b \cup_5 c)$, $b \cup (a \cup_5 c)$, and $c \cup (a \cup_5 b)$, respectively.

Let us now consider the case $i = 3$, Eq. (5.1). According to the first definition of aCb from Definition 2.3 we have, given the premises aCb and aCc and the orthomodularity property [$a \cup (a' \cap (a \cup b)) = a \cup b$]:

$$a \cup_3 b = (a \cap b) \cup (a \cap b') \cup (a' \cap (a \cup b)) = a \cup (a' \cap (a \cup b)) = a \cup b$$

and therefore, using F–H theorem and the second premise and Definition 2.3,

$$\begin{aligned} (a \cup_3 b) \cup_3 c &= ((a \cup b) \cap c) \cup ((a \cup b) \cap c') \cup ((a' \cap b') \cap (a \cup b \cup c)) \\ &= [\text{F–H}] \\ &= (a \cap c) \cup (b \cap c) \cup (a \cap c') \cup (b \cap c') \cup ((a' \cap b') \cap (a \cup b \cup c)) \\ &= [\text{Def. 2.3}] = a \cup (b \cap c) \cup (b \cap c') \cup ((a' \cap b') \cap (a \cup b \cup c)) \\ &= (b \cap c) \cup (b \cap c') \cup ((a \cup (a' \cap b')) \cap (a \cup b \cup c)) \\ &= (b \cap c) \cup (b \cap c') \cup (((a \cup a') \cap (a \cup b')) \cap (a \cup b \cup c)) \\ &= (b \cap c) \cup (b \cap c') \cup ((a \cup b') \cap (a \cup b \cup c)) \end{aligned} \quad (5.7)$$

The right-hand side of the conclusion in Eq. (5.1) reads

$$a \cup_3 (b \cup_3 c) = (a \cap (b \cup_3 c)) \cup (a \cap (b \cup_3 c)') \cup (a' \cap (a \cup (b \cup_3 c))). \quad (5.8)$$

Now $b \cup_3 c = (b \cap c) \cup (b \cap c') \cup (b' \cap (b \cup c))$ and since we also have aCb and aCc and therefore, $aC(b \cap c)$, $aC(b \cap c')$, and $aC(b' \cap (b \cup c))$; we have $aC(b \cup_3 c)$ as well. Hence, using Definition 2.3 we reduce Eq. (5.8) to

$$\begin{aligned} a \cup_3 (b \cup_3 c) &= a \cup a' \cap (a \cup (b \cup_3 c)) = a \cup (b \cap c) \cup (b \cap c') \cup (b' \cap (b \cup c)) \\ &= (b \cap c) \cup (b \cap c') \cup ((a \cup b') \cap (a \cup b \cup c)), \end{aligned}$$

which is nothing but Eq. (5.7). Hence, Eq. (5.1) is proved.

Let us next consider Eq. (5.2). Here we have, $a \cup_3 b = a \cup b$ and $b \cup_3 c = b \cup c$ and therefore,

$$\begin{aligned}
 (a \cup_3 b) \cup_3 c &= ((a \cup b) \cap c) \cup ((a \cup b) \cap c') \cup ((a' \cap b') \cap (a \cup b \cup c)) \\
 &= [\text{F-H}] \\
 &= (a \cap c) \cup (b \cap c) \cup (a \cap c') \cup (b \cap c') \cup ((a' \cap b') \cap (a \cup c)) \\
 &= (a \cap c) \cup (a \cap c') \cup (b \cap c) \cup (b \cap c') \cup ((a' \cap b') \cap (a \cup c)) \\
 &= [bCc] = (a \cap c) \cup (a \cap c') \cup b \cup ((a' \cap b') \cap (a \cup c)) \\
 &= [bC(a' \cap b'), bC(b \cup c)] \\
 &= (a \cap c) \cup (a \cap c') \cup ((a' \cup b) \cap (a \cup b \cup c)). \tag{5.9}
 \end{aligned}$$

On the other hand, we have

$$\begin{aligned}
 a \cup_3 (b \cup_3 c) &= (a \cap (b \cup c)) \cup (a \cap b' \cap c') \cup (a' \cap (a \cup b \cup c)) \\
 &= [\text{F-H}] = (a \cap b) \cup (a \cap c) \cup (a \cap b' \cap c') \cup (a' \cap (a \cup b \cup c)) \\
 &= (a \cap c) \cup (a \cap b' \cap c') \cup (a \cap b) \cup (a' \cap b) \cup (a' \cap (a \cup c)) \\
 &= [aCb] = (a \cap c) \cup (a \cap b' \cap c') \cup b \cup (a' \cap (a \cup c)) \\
 &= [bC(a \cap c')] \\
 &= (a \cap c) \cup ((a \cap c') \cup b) \cap (b' \cup b) \cup (a' \cap (a \cup c)) \\
 &= (a \cap c) \cup (a \cap c') \cup b \cup (a' \cap (a \cup c)) \\
 &= [bCa', bC(a \cup c)] = (a \cap c) \cup (a \cap c') \cup ((a' \cup b) \cap (a \cup b \cup c))
 \end{aligned}$$

which is nothing but Eq. (5.9) and this proves Eq. (5.2).

As for Eq. (5.3), here we again have $cC(a \cup_3 b)$ and $b \cup_3 c = b \cup c$. Thus we get,

$$\begin{aligned}
 (a \cup_3 b) \cup_3 c &= ((a \cup_3 b) \cap c) \cup ((a \cup_3 b) \cap c') \cup ((a \cup_3 b)' \cap ((a \cup_3 b) \cup c)) \\
 &= (a \cup_3 b) \cup ((a \cup_3 b)' \cap ((a \cup_3 b) \cup c)) \\
 &= [\text{OM property}] = (a \cup_3 b) \cup c \\
 &= (a \cap b) \cup (a \cap b') \cup (a' \cap (a \cup b)) \cup c \\
 &= (a \cap b) \cup (a \cap b') \cup ((a' \cup c) \cap (a \cup b \cup c)) \tag{5.10}
 \end{aligned}$$

For the right-hand side we have,

$$\begin{aligned}
 a \cup_3 (b \cup_3 c) &= (a \cap (b \cup c)) \cup (a \cap b' \cap c') \cup (a' \cap (a \cup b \cup c)) \\
 &= (a \cap b) \cup (a \cap c) \cup (a \cap b' \cap c') \cup (a' \cap (a \cup b \cup c))
 \end{aligned}$$

$$\begin{aligned}
 &= (a \cap b) \cup (a \cap b' \cap c') \cup (a' \cup (a \cap c)) \cap ((a \cap c) \cup a \cup b \cup c) \\
 &= (a \cap b) \cup (a \cap b' \cap c') \cup (a' \cup c) \cap (a \cup b \cup c) \\
 &= (a \cap b) \cup (a \cap b' \cap c') \cup (a' \cap (a \cup b \cup c)) \cup c \\
 &= (a \cap b) \cup ((a \cap b') \cup c) \cap (c \cup c') \cup (a' \cap (a \cup b \cup c)) \\
 &= (a \cap b) \cup (a \cap b') \cup c \cup (a' \cap (a \cup b \cup c)) \\
 &= (a \cap b) \cup (a \cap b') \cup ((a' \cup c) \cap (a \cup b \cup c))
 \end{aligned}$$

which is nothing but Eq. (5.10), which proves Eq. (5.3).

Since $a \cup_4 b = b \cup_3 a$ and $a \cap_{3,4} b = (a' \cup_{3,4} b)'$, we have proved the theorem for $i = 3, 4$. \square

We conjecture that the theorem holds in any weakly orthomodular lattice, WOML (Pavičić and Megill, 1998a) as well.

6. CONDITIONAL DISTRIBUTIVITY OF QUANTUM OPERATIONS

The F–H theorem for conditional distributivity does not in general hold for the quantum disjunctions and conjunctions. D’Hooghe and Pykacz show this for \cup_5, \cap_5 [2000, p. 646] and state (in our notation for i) “the same can be checked for $i = 1, 2, 3, 4$ ” (p. 647). While this is true for $i = 3, 4$, distributivity in the forms given by Theorems 6.1 and 6.2 does hold for $i = 1, 2$. Also, parts of the Foulis–Holland theorem, presented in Theorem 6.3 hold for any i and therefore for the unified quantum disjunction and conjunction (\cup and \cap) from section 4.

Theorem 6.1. *In any orthomodular lattice any triple $\{a, b, c\}$ in which one of the elements commutes with the other two is distributive with respect to \cup_1 and \cap_1 in the following sense:*

$$aCb \ \& \ aCc \ \Rightarrow \ a \cup_1 (b \cap_1 c) = (a \cup_1 b) \cap_1 (a \cup_1 c) \tag{6.1}$$

$$aCb \ \& \ bCc \ \Rightarrow \ a \cup_1 (b \cap_1 c) = (a \cup_1 b) \cap_1 (a \cup_1 c) \tag{6.2}$$

$$aCc \ \& \ bCc \ \Rightarrow \ a \cup_1 (b \cap_1 c) = (a \cup_1 b) \cap_1 (a \cup_1 c) \tag{6.3}$$

Proof. In this and all other proofs of this section, we will implicitly make use of the rules $aCb \Rightarrow a \cup_i b = a \cup_j b$, $aCb \Rightarrow a \cap_i b = a \cap_j b$, $aCb \ \& \ aCc \Rightarrow aCb \cup_i \cap_j c$, and $aCb \Rightarrow a, b, a \cup_i b, a \cap_i b C a \cup_j b, a \cap_j b, 0 \leq i, j \leq 5$. Also, $aCa \cup_{0,1,3,5}, \cap_{0,1,3,5} b, b C a \cup_{0,2,4,5}, \cap_{0,2,4,5} b, a \cup_{0,1} b C a' \cup_{0,1} c, a \cap_{0,1} b C a' \cap_{0,1} c, a \cup_{0,1} b C c \cup_{0,2} a',$ and $a \cap_{0,1} b C c \cap_{0,2} a'$. We will use F–H implicitly.

Recall that $\cup_0 = \cup$.

$$\begin{aligned} \text{For (6.1), } a \cup (b \cap_1 c) &= a \cup (b \cap (b' \cup c)) = (a \cup b) \cap ((a' \cap b') \cup a \cup c) \\ &= (a \cup b) \cap_1 (a \cup c). \end{aligned}$$

$$\begin{aligned} \text{For (6.2), } a \cup_1 (b \cap c) &= a \cup (a' \cap b \cap c) = a \cup (b \cap a' \cap c) \\ &= (a \cup b) \cap (a \cup (a' \cap c)) = (a \cup b) \cap (a \cup_1 c). \end{aligned}$$

$$\begin{aligned} \text{For (6.3), } a \cup_1 (b \cap c) &= a \cup (a' \cap b \cap c) = (a \cup (a' \cap b)) \cap (a \cap c) \\ &= (a \cup_1 b) \cap (a \cup c). \quad \square \end{aligned}$$

Because \cup_1, \cap_1 are not commutative, the “reverse” distributivity $(a \cap_1 b) \cup_1 c = (a \cup_1 c) \cap_1 (b \cup_1 c)$ does not hold for all F–H hypotheses. However, it does hold for \cup_2, \cap_2 :

Theorem 6.2. *In any orthomodular lattice any triple $\{a, b, c\}$ in which one of the elements commutes with the other two is distributive with respect to \cup_2 and \cap_2 in the following sense:*

$$aCb \ \& \ aCc \ \Rightarrow \ (a \cap_2 b) \cup_2 c = (a \cup_2 c) \cap_2 (b \cup_2 c) \tag{6.4}$$

$$aCb \ \& \ bCc \ \Rightarrow \ (a \cap_2 b) \cup_2 c = (a \cup_2 c) \cap_2 (b \cup_2 c) \tag{6.5}$$

$$aCc \ \& \ bCc \ \Rightarrow \ (a \cap_2 b) \cup_2 c = (a \cup_2 c) \cap_2 (b \cup_2 c) \tag{6.6}$$

Proof. Theorem 6.1 and the fact that $a \cup_2 b = b \cup_1 a, a \cap_2 b = b \cap_1 a$. \square

For certain F–H hypotheses, distributive laws hold for all $i = 1, \dots, 5$. In addition, a couple of other cases hold for $i = 1, 2$.

Theorem 6.3. *In any orthomodular lattice the following laws hold:*

$$aCb \ \& \ aCc \ \Rightarrow \ a \cup_i (b \cap_i c) = (a \cup_i b) \cap_i (a \cup_i c), \quad i = 1, \dots, 5 \tag{6.7}$$

$$aCc \ \& \ bCc \ \Rightarrow \ (a \cap_i b) \cup_i c = (a \cup_i c) \cap_i (b \cup_i c), \quad i = 1, \dots, 5 \tag{6.8}$$

$$aCb \ \& \ aCc \ \Rightarrow \ (a \cap_1 b) \cup_1 c = (a \cup_1 c) \cap_1 (b \cup_1 c) \tag{6.9}$$

$$aCc \ \& \ bCc \ \Rightarrow \ a \cup_2 (b \cap_2 c) = (a \cup_2 b) \cap_2 (a \cup_2 c) \tag{6.10}$$

Proof. For Eq. (6.7), using $aCb \Rightarrow a \cup_i b = a \cup b$ and $aCb \ \& \ aCc \Rightarrow aC(b \cup_i c)$ and F–H we can write the conclusion as

$$a \cup (b \cap_i c) = (a \cup b) \cap_i (a \cup c), \quad i = 1, \dots, 5 \tag{6.11}$$

To prove that the right-hand side boils down to the left-hand one is straightforward and can be done in a complete analogy to the case $i = 1$ already done previously – Eq. (6.1). For example, for $i = 4$ we have,

$$\begin{aligned} (a \cup b) \cap_4 (a \cup c) &= ((a' \cap c') \cup ((a \cup b) \cap (a \cup c))) \cap (a \cup b \cup c) \\ &\quad \cap ((a' \cap b') \cup a \cup c) \\ &= ((a' \cap c') \cup a \cup (b \cap c)) \cap (a \cup b \cup c) \cap (a \cup b' \cap c) \\ &= a \cup (b \cap_4 c). \end{aligned}$$

For Eq. (6.8), the proof follows from Eq. (6.11) by symmetry.

The proof of (6.9) seems a little tricky, so we show it in some detail. First, we show that (under the hypotheses)

$$(a \cap b) \cup (a' \cap c) = (a \cup c) \cap (b \cup a'). \tag{6.12}$$

From $b \geq a \cap b = a \cap (b \cup a')$ and $c \geq c \cap (b \cup a')$ we have $b \cup c \geq (a \cap (b \cup a')) \cup (c \cap (b \cup a')) = (a \cup c) \cap (b \cup a')$. Therefore $(a \cup c) \cap (b \cup a') = (b \cup a') \cap (a \cup c) \cap (b \cup c) = ((a \cap b) \cup a') \cap ((a \cap b) \cup c) = (a \cap b) \cup (a' \cap c)$, establishing (6.12). The left-hand side of (6.9) reduces to $(a \cap b) \cup_1 c = (a \cap b) \cup ((a \cap b)' \cap c) = (a \cap b) \cup ((a' \cup b') \cap c) = (a \cap b) \cup (a' \cap c) \cup (b' \cap c)$. The right-hand side reduces to $(a \cup c) \cap_1 (b \cup_1 c) = (a \cup c) \cap ((a \cup c)' \cup b \cup (b' \cap c)) = (a \cup c) \cap ((a' \cap c') \cup b \cup (b' \cap c)) = (a \cup c) \cap ((b \cup a') \cap (b \cup c')) \cup (b' \cap c) = (a \cup c) \cap ((b' \cap c) \cup_1 (b \cup a')) = (a \cup c) \cap ((b \cup a') \cup (b' \cap c)) = ((a \cup c) \cap (b \cup a')) \cup ((a \cup c) \cap b' \cap c) = ((a \cup c) \cap (b \cup a')) \cup (b' \cap c)$. Using (6.12), we see they are the same.

For (6.10) we use (6.9) and $a \cup_2 b = b \cup_1 a, a \cap_2 b = b \cap_1 a$. \square

Similar results can be stated for the dual operations (\cup_i and \cap_i interchanged). In all other cases not shown in the three theorems above, the distributive law does not hold: all of them fail in orthomodular lattice MO2 (Fig. 1).

If we allow a mixture of the different disjunctions and conjunctions, we can obtain a distributive law that holds unconditionally.

Theorem 6.4. *In any orthomodular lattice the following law holds:*

$$a \cup_1 (b \cap_0 c) = (a \cup_1 b) \cap_0 (a \cup_1 c) \tag{6.13}$$

Proof. Expanding definitions and using F–H, $a \cup_1 (b \cap_0 c) = a \cup (a' \cap b \cap c) = a \cup (a' \cap b \cap a' \cap c) = (a \cup (a' \cap b)) \cap (a \cup (a' \cap c)) = (a \cup_1 b) \cap_0 (a \cup_1 c)$. \square

It is interesting that if we consider all equations of the form $a \cup_i (b \cap_j c) = (a \cup_k b) \cap_l (a \cup_m c)$ for all possible assignments $0 \leq i, j, k, l, m \leq 5$ ($6^5 = 7776$




Fig. 1. Lattice MO2.

possibilities), the equation holds in all OMLs for exactly the one case of (6.13): $i = 1, j = 0, k = 1, l = 0, m = 1$. All other 7775 cases fail in lattice MO2.

The “reverse” form of (6.13) holds with \cup_2 substituted for \cup_1 . Dual results with \cup_i and \cap_i interchanged can also be stated.

7. AN OPEN PROBLEM

In Megill and Pavičić (2000) we opened an interesting problem on whether the “distributivity of symmetric identity,” expressed by Eq. (7.7), holds in all orthomodular lattices or not and if a particular equation derivable from it in any orthomodular lattice characterizes the latter lattices. An indication that they might do so is that they pass all Greechie diagrams we let them run on – with up to 38 atoms and 38 blocks (more than 50 million lattices). We used our program *greechie* to obtain the diagrams and our program *latticeg* to check the equations on them (Mckay *et al.*, 2000). On the other hand Eq. (7.7) does not imply the orthomodularity property – it does not fail in the diagram 06, which characterizes all orthomodular lattices.

In Megill and Pavičić (2000) we proved several partial results for the above distributivity. In this section we prove that it holds in Hilbert space and in the Godowski lattices of the second lowest order (4GO). We recall from Megill and Pavičić (2000) that a 4GO is any OML (actually any OL) in which the following equation, which we call 4-Go, holds:

$$(a \rightarrow_1 b) \cap (b \rightarrow_1 c) \cap (c \rightarrow_1 d) \cap (d \rightarrow_1 a) \leq a \rightarrow_1 d. \tag{7.1}$$

We define $a \equiv b \stackrel{\text{def}}{=} (a \cap b) \cup (a' \cap b')$ and note that $a \equiv b = a \equiv_5 b$ holds in all OMLs.

Lemma 7.1. *In any OML we have,*

$$(a \equiv c) \cup (b \equiv c) = ((a \rightarrow_2 c) \cup (b \rightarrow_2 c)) \cap ((c \rightarrow_1 a) \cup (c \rightarrow_1 b)) \quad (7.2)$$

$$(a \equiv c) \cup (b \equiv c) \leq ((a \cap b) \rightarrow_2 c) \cap (c \rightarrow_1 (a \cup b)) \quad (7.3)$$

$$((a \cup b) \equiv c) \cap (a \equiv b) = (a \equiv c) \cap (a \equiv b). \quad (7.4)$$

In any 4GO we have,

$$(a \equiv b) \cap ((b \equiv c) \cup (a \equiv c)) \leq a \equiv c. \quad (7.5)$$

Proof. For (7.2), we have,

$$\begin{aligned} (a \equiv c) \cup (b \equiv c) &= (b \cap c) \cup (a' \cap c') \cup (b' \cap c') \cup (a \cap c) \\ &= (b \cap c) \cup (a' \cap c') \cup ((b \rightarrow_2 c) \cap (c \rightarrow_1 a)) \\ &= (b \cap c) \cup (((a' \cap c') \cup (b \rightarrow_2 c)) \cap ((a' \cap c') \cup (c \rightarrow_1 a))) \\ &= (b \cap c) \cup (((a' \cap c') \cup (b \rightarrow_2 c)) \cap (c \rightarrow_1 a)) \\ &= ((b \cap c) \cup (a' \cap c') \cup (b \rightarrow_2 c)) \cap ((b \cap c) \cup (c \rightarrow_1 a)) \\ &= ((a' \cap c') \cup (b \rightarrow_2 c)) \cap ((b \cap c) \cup (c \rightarrow_1 a)) \\ &= (((a' \cap c') \cup c) \cup (c \cup (b' \cap c'))) \cap (((b \cap c) \cup c') \\ &\quad \cup (c' \cup (c \cap a))) \\ &= ((a \rightarrow_2 c) \cup (b \rightarrow_2 c)) \cap ((c \rightarrow_1 b) \cup (c \rightarrow_1 a)). \end{aligned}$$

In the second step, we use Eq. (3.20) from Megill and Pavičić (2000). In the third and fifth steps we apply the F–H theorem, and in the fourth and sixth steps we apply absorption laws.

For (7.3), $(a \rightarrow_2 c) \cup (b \rightarrow_2 c) \leq (a \cap b) \rightarrow_2 c$ and $(c \rightarrow_1 a) \cup (c \rightarrow_1 b) \leq c \rightarrow_1 (a \cup b)$ in any OL, so $(a \equiv c) \cup (b \equiv c) = [\text{from (7.2)}]((a \rightarrow_2 c) \cup (b \rightarrow_2 c)) \cap ((c \rightarrow_1 a) \cup (c \rightarrow_1 b)) \leq ((a \cap b) \rightarrow_2 c) \cap (c \rightarrow_1 (a \cup b))$.

For (7.4), we have,

$$\begin{aligned} ((a \cup b) \equiv c) \cap (a \equiv b) &= ((a \cup b) \cap c) \cup (a' \cap b' \cap c') \cap ((a \cap b) \cup (a' \cap b')) \\ &= (((a \cup b) \cap c) \cap ((a \cap b) \cup (a' \cap b'))) \cup ((a' \cap b' \cap c') \\ &\quad \cap ((a \cap b) \cup (a' \cap b'))) \\ &= (((a \cup b) \cap c) \cap (a \cap b)) \cup (((a \cup b) \cap c) \cap (a' \cap b')) \\ &\quad \cup ((a' \cap b' \cap c') \cap (a \cap b)) \cup ((a' \cap b' \cap c') \cap (a' \cap b')) \\ &= ((a \cap b \cap c) \cup 0 \cup 0 \cup (a' \cap b' \cap c')) \\ &= (a \equiv c) \cap (a \equiv b) \end{aligned}$$

where in the second and third steps we apply F–H and in the last step we apply Lemma 3.11 of Megill and Pavičić (2000).

Finally, (7.5) is proved as follows. Equation (3.30) of Megill and Pavičić (2000), which we repeat below as (7.6), was shown to hold in all 4GOs.

$$(a \equiv b) \cap ((b' \cap c') \cup (a \cap c)) \leq a \equiv c. \tag{7.6}$$

Using Eq. (3.20) of Megill and Pavičić (2000) and renaming variables, we see that this is the same as

$$(d \equiv e) \cap (e \rightarrow_2 c) \cap (c \rightarrow_1 d) \leq d \equiv c.$$

Substituting $a \cup b$ for d and $a \cap b$ for e ,

$$((a \cup b) \equiv (a \cap b)) \cap ((a \cap b) \rightarrow_2 c) \cap (c \rightarrow_1 (a \cup b)) \leq (a \cup b) \equiv c.$$

Since $(a \cup b) \equiv (a \cap b) = a \equiv b$ holds in any OML, we have

$$\begin{aligned} (a \equiv b) \cap ((a \cap b) \rightarrow_2 c) \cap (c \rightarrow_1 (a \cup b)) &\leq (a \cup b) \equiv c \\ (a \equiv b) \cap ((a \cap b) \rightarrow_2 c) \cap (c \rightarrow_1 (a \cup b)) &\leq ((a \cup b) \equiv c) \cap (a \equiv b) \\ (a \equiv b) \cap ((a \cap b) \rightarrow_2 c) \cap (c \rightarrow_1 (a \cup b)) &\leq (a \equiv c) \cap (a \equiv b) \\ (a \equiv b) \cap ((a \cap b) \rightarrow_2 c) \cap (c \rightarrow_1 (a \cup b)) &\leq a \equiv c \\ (a \equiv b) \cap ((a \equiv c) \cup (b \equiv c)) &\leq a \equiv c. \end{aligned}$$

where in the third step we use (7.4) and in the last step we use (7.3). \square

Theorem 7.2. *The following equation, which we call distributivity of symmetric identity, holds in all 4GOs (and therefore all nGO, $n \geq 4$) and thus in the lattice of all closed subspaces of finite- or infinite-dimensional Hilbert space:*

$$(a \equiv b) \cap ((b \equiv c) \cup (a \equiv c)) = ((a \equiv b) \cap (b \equiv c)) \cup ((a \equiv b) \cap (a \equiv c)). \tag{7.7}$$

Proof. The result follows immediately from (7.5) and Theorem 2.9 of Megill and Pavičić (2000). \square

Whether or not (7.7) holds in all OMLs or even in all WOMLs (since it does not fail in O6) is still an open question. However, the most important question from the point of view of quantum mechanics, which is whether or not it holds in Hilbert space, is answered by Theorem 7.2.

Since (7.6) also follows from (7.7), as shown in Megill and Pavičić (2000), the OML variety in which (7.6) holds is the same as the OML variety in which (7.7) holds. Thus if one of these holds in any OML (our open question), so does the other.

Another open question is whether the stronger-looking Eq. (3.29) of Megill and Pavičić (2000), from which (7.6) follows and which we repeat here as (7.8),

$$(a \rightarrow_1 b) \cap (b \rightarrow_2 c) \cap (c \rightarrow_1 a) \leq (a \equiv c) \quad (7.8)$$

can be derived (in an OML) from (7.6).

8. ALGORITHMS FOR THE PROGRAMS

In an OML, any expression with two variables is equal to 1 of 96 canonical forms, corresponding to the 96 elements of the free OML F_2 . We fix the 96 expressions of Beran (1985, Table 1, p. 82) as our canonical standard.

The program `beran` takes, as its input, an arbitrary two-variable expression and outputs the equivalent canonical form. The program can be used to prove or disprove any two-variable conjecture expressed as an equation, simply by verifying that both sides of the equation reduce (or do not reduce) to the same canonical form.

Each element of OML F_2 can be separated into a “Boolean part” and an “MO2 part” (Navara, 1997). Each of them has relatively simple rules of calculation, and we use this method in the program `beran`.⁶ This is implemented in the program by checking for either Boolean or MO2 lattice violation of the 96 equations formed by setting the input expression equal to each of the 96 canonical expressions, and the unique equation that violates neither lattice gives us the answer.

The program `beran` is contained in a single file, `beran.c`, and compiles on any platform with an ANSI C compiler such as `gcc`. The use of the program is simple. The operations \cup , \cap , and $'$ are represented by the characters `v`, `^`, and `-`. (Other operations are also defined and can be seen with the program’s `—help` option). As an example, to see the canonical expression corresponding to $a \cup (a' \cap (a \cup b))$, we type

```
beran "(av(-a^(avb)))"
```

and the program responds with `(avb)`.

A second program, `bercomb`, was used to find the minimal expressions shown in section 3. This program is contained in the single file `bercomb.c`. Its input parameters include the number of variable occurrences and the number of negations (orthocomplementations), and it exhaustively constructs all possible expressions containing a single binary operation with these parameters fixed. For each expression it uses the algorithm from `beran.c` to determine the expression’s canonical form which it prints out. When a set of operations is specified, such as \rightarrow_1 through \rightarrow_5 , it prints out the canonical form only when all operations simultaneously result in that canonical form.

⁶The authors wish to thank Prof. Navara for suggesting this method. The reader can download this or any other afore-mentioned program from our ftp sites: <ftp://m3k.grad.hr/pavicic/quantum-logic/programs/> and <ftp://users.shore.net/members/n/d/ndm/quantum-logic/>

If $v \geq 2$ is the number of variable occurrences and n is ≥ 0 and $\leq (2v - 1)$ the number of negations, the number of possible expressions containing one or two different variables is as follows. The number of ways of parenthesizing a binary operation in an expression with v variables is the Catalan number C_{v-1} , where $C_i = \binom{2i}{i}/(i + 1)$. There are 2^v possible ways to assign two variables to an expression. If we display an expression with no negations in Polish notation, it is easy to see that there are $2v - 1$ symbols and therefore $\binom{2v-1}{n}$ ways to distribute n negations (disallowing double negations). Thus for fixed v and n , there are $2^v \binom{2v-1}{n} C_{v-1}$ possible expressions.

For example, if we type

```
bercomb 7 0 i n
```

then all $2^7 \binom{2 \cdot 7 - 1}{0} C_{7-1} = 16896$ expressions with 7 variable occurrences and 0 negations are scanned, and the output includes the four smallest implicational expressions resulting in $a \cup b$ that we mentioned before Lemma 3.1. We refer the reader to the program's `—help` option for the meaning and usage of the other `bercomb` parameters. In this example `i` means \rightarrow_1 through \rightarrow_5 and `n` means don't suppress duplicate canonical expressions.

9. CONCLUDING REMARKS

In Pavičić and Megill (1999) we stressed that all the operations in an orthomodular lattice are fivefold defined and we illustrated this on the identity operations. The claim was based on Pavičić and Megill (1998b) where we proved that “quantum” as well as “classical” operations can serve for a formulation of an orthomodular lattice underlying Hilbert space. In 1998, we also put on the web the computer program `beran`, which reduces any two-variable expression in an orthomodular lattice to one of the 96 possible ones as given in Beran (1985).

In effect, in the standard orthomodular lattice formulation (where the “classical” operations are inherited from the Hilbert space formalism) there are 80 quantum expressions that for compatible variables reduce to 16 classical expressions. In general all quantum expressions (including “quantum 0” and “quantum 1”) are fivefold defined. (Detailed presentation of them all we give in section 2.) In our quantum algebraic approach the situation reverses and we have classical operations fivefold defined in a quantum algebra formulation.

Still, recently several papers on “some new operations on orthomodular lattices” appeared in press as, for example, the one by D’Hooghe and Pykacz (2000) in which they picked out Beran expressions 12, 18, 28, 34, 44, 50, 60, and 76 and looked at some of their properties. So, for example, in D’Hooghe and Pykacz (2000, p. 649, bottom) one reads (in our notation): “Theorem 7 allows one to express in many ways any of the studied operations by (any of) the other(s) orthocomplementation. However, the following example in which we

express \cup_1 by \cup_5 and shows that the obtained formulas might be rather lengthy: $a \cup_1 b = (a \cup_5 ((a \cup_5 b)' \cup_5 a))' \cup_5 a$. It is an open question which of such formulas (if any) could be written in a more economical way.” Our approach immediately closes this open question: all the formulas could be written in a more economical way and one gets all alternatives in seconds; for example, in the afore-cited example, there are over 100 shorter expressions—one of 3 shortest ones is given by Eq. (3.16)—and there are over 500 of them with the same (5) variable occurrences. On the other hand, Theorem 6 from D’Hooghe and Pykacz (2000, p. 648) is just a special case of our Theorem 2.5 from Pavičić (1993, p. 1487). Also all the results from D’Hooghe and Pykacz (2000, section 3.2, p. 646–648), can be trivially obtained using our computer program `latticeg` (McKay *et al.*, 2000). In addition, their two conjectures (p. 648) following from their Theorem 5 (p. 647) one can support by our program `latticeg` with millions of lattices. Hence, it appears necessary to present our results in detail, give explicit proofs of all our previous claims, present the most important and relevant outputs of our programs in some detail, and provide the reader with instructions on how to use our programs that give answers to virtually all questions one can have on algebraic properties of two-variable orthomodular formulas in seconds.

Thus, in section 3 we prove several lemmas in which we show how one can express operations in any standardly defined (in section 2) orthomodular lattice by each other. Lemma 3.2 gives expressions of classical disjunction (\cup) by means of all five quantum implications \rightarrow_i , $i = 1, \dots, 5$ and without negations in a shortest possible single equation—meaning that the equation preserves its form for all i ’s and that there are no simpler equations with such a property. Expressions of \cup by means of quantum disjunctions (\cup_i , $i = 1, \dots, 5$) and conjunctions (\cap_i , $i = 1, \dots, 5$) follow from Definition 2.4. Lemma 3.2 gives the shortest expressions of $\cup = \cup_0$ and $\cap = \cap_0$ by means of \rightarrow_i , \cup_i , and \cap_i , $i = 1, \dots, 5$, with negation. Lemma 3.4 gives the shortest expressions of \cup_i , \cap_i , and \equiv_i , $i = 0, \dots, 5$, by means of \cup_i and \cap_i , $i = 1, \dots, 5$, with negation.

In section 4 we start with the possibility—opened by Lemma 3.2—of expressing \cup by means of \cup_i , $i = 1, \dots, 5$, in five equations of the same form and define—in Definition 4.1—the orthomodular lattice by means of one unique quantum operation. We have chosen quantum disjunction \mathbb{U} , but the same, of course, can be done with quantum conjunction \mathbb{M} or implication (the latter being just another way of writing disjunction) – quantum identity is the only quantum operation that cannot serve the purpose, as we proved in Pavičić (1998). In such a formulation of orthomodular lattice everything reverses and now classical operations can be expressed in five different ways as shown by Theorem 4.3.

We stress that the quantum algebra QA (Definition 4.1) is actually completely defined by its Substitution Rule, and that “axioms” A1–A7 are merely some consequences of that rule. A1–A7 are important in that they show that standard OML

can be embedded in QA and are included for this reason. However there are many other nonobvious consequences of QA such as those exemplified in Lemma 4.2. That lemma only touches the surface of the kind of conditions one can obtain from QA, and it is possible that QA provides a rich algebraic structure that has yet to be explored. It also remains an interesting open problem if QA can be finitely axiomatized.

Lemma 3.3 shows the surprising result that classical disjunction can be expressed in a single equation that holds in any OML for *all* 6 disjunctions $\cup_i, i = 0, \dots, 5$. This opens the possibility of an even more general quantum algebra, with Eq. (3.7) used in place Eq. (3.4) as the basis for A1–A7. In this case we would replace “ $i = 0, \dots, 5$ ” for “ $i = 1, \dots, 5$ ” in the Substitution Rule. The same kinds of open questions we brought up for QA would also apply to this more general algebra.

As for D’Hooghe and Pykacz’s conjecture on a possible conditional associativity of $\cup_{3,4}$ and $\cap_{3,4}$ (Hooghe and Pykacz, 2000), we decided that its passage through millions of Greechie diagrams makes it worth proving and we did so in section 5. In that way we obtained the conditional associativity for the unified operations \cup and \cap from section 5.

In section 6 we prove several Foulis–Holland-type conditional distributivities, some of which are valid for all standard quantum disjunctions and conjunctions and therefore for the unified quantum disjunction and conjunction (\cup and \cap) from section 4.

As for properties taken over from Hilbert space in section 7 we present two, given by Eqs. (7.5) and (7.7), which proved to hold in a variety of orthomodular lattice 4GO (and therefore in n GO, $n \geq 4$), but which do not fail in any of over 50 million Greechie diagrams we tested the properties on. Thus it remains an open problem whether the properties hold in any orthomodular lattice and even more whether Eq. (7.7) holds in an even weaker ortholattice called weakly orthomodular lattice, WOML.

In section 8 we give algorithms we used to obtain and check all our equations and proofs for properties involving two variables.

To conclude, the only genuine target that apparently remains for scientific investigation in algebraic properties of orthomodular lattices in the future are properties with three and more variables.

ACKNOWLEDGMENTS

One of us (M. P.) is grateful to Prof. Anton Zeilinger, Institute for Experimental Physics, University of Vienna, Austria, for his kind invitation to participate in the program *Quantum Measurement Theory and Quantum Information* (Sept. 1, 2000–Jan. 20, 2001) organized by A. Eckert, A. Zeilinger and P. Zoller at the Erwin

Schrödinger International Institute for Mathematical Physics in Vienna, where this paper has been written in part. He would also like to acknowledge support of the Ministry of Science of Croatia through the Project No. 082006.

REFERENCES

- Beran, L. (1985). *Orthomodular Lattices; Algebraic Approach*, D. Reidel, Dordrecht.
- Dalla Chiara, M. L. (1986). Quantum logic. In *Handbook of Philosophical Logic*, D. Gabbay and F. Guenther, eds., D. Reidel, Dordrecht, Vol. 3, pp. 427–469.
- Dalla Chiara, M. L. and Giuntini, R. (2001). Quantum logics. <http://xxx.lanl.gov/abs/quant-ph/0101028>.
- D’Hooghe, B. and Pykacz, J. (2000). On some new operations on orthomodular lattices. *International Journal of Theoretical Physics* **39**, 641.
- Dishkant, H. (1974). The first order predicate calculus based on the minimal logic of quantum mechanics. *Report on Mathematics Logic* **3**.
- Halmos, P. R. (1957). *Introduction to Hilbert Space and the Spectral Theory of Spectral Multiplicity*, Chelsea, New York.
- Holland, S. S., JR (1995). Orthomodularity in infinite dimensions; a theorem of M. Solèr. *Bulletin of the American Mathematics Society* **32**, 205.
- Isham, C. J. (1995). *Lectures of Quantum Theory*, Imperial College Press, London.
- Kalmbach, G. (1974). Orthomodular logic. *Zeitschrift für mathematische Logik und Grundlagen der Mathematik* **20**, 395.
- Kalmbach, G. (1983). *Orthomodular Lattices*, Academic Press, London.
- McKay, B. D., Megill, N. D., and Pavičić, M. (2000). Algorithms for Greechie diagrams. *International Journal of Theoretical Physics* **39**, 2393.
- Megill, N. D. and Pavičić, M. (2000). Equations, states, and lattices of infinite-dimensional Hilbert space. *International Journal of Theoretical Physics* **39**, 2349.
- Mittelstaedt, P. (1978). *Synthese Library, Vol. 18: Quantum Logic*, Reidel, London.
- Navara, M. (1997). On generating finite orthomodular sublattices. *Tatra Mountains Mathematics Publications* **10**, 109.
- Pavičić, M. (1987). Minimal quantum logic with merged implications. *International Journal of Theoretical Physics* **26**, 845.
- Pavičić, M. (1989). Unified quantum logic. *Foundation Physics* **19**, 999–1016.
- Pavičić, M. (1993). Nonordered quantum logic and its YES-NO representation. *International Journal of Theoretical Physics* **32**, 1481.
- Pavičić, M. (1998). Identity rule for classical and quantum theories. *International Journal of Theoretical Physics* **37**, 2099.
- Pavičić, M. and Megill, N. D. (1998a). Binary orthologic with modus ponens is either orthomodular or distributive. *Helvetica Physica Acta* **71**, 610.
- Pavičić, M. and Megill, N. D. (1998b). Quantum and classical implication algebras with primitive implications. *International Journal of Theoretical Physics* **37**, 2091.
- Pavičić, M. and Megill, N. D. (1999). Non-orthomodular models for both standard quantum logic and standard classical logic: Repercussions for quantum computes. *Helvetica Physica Acta* **72**, 189, 1999. <http://xxx.lanl.gov/abs/quant-ph/9906101>.
- Zeman, J. J. (1979). Quantum logic with implications. *Notre Dame Journal of Formal Logic* **20**, 723.