

Quantum and Classical Implication Algebras with Primitive Implications

Mladen Pavičić¹ and Norman D. Megill²

Received November 30, 1997

Join in an orthomodular lattice is obtained in the same form for all five quantum implications. The form holds for the classical implication in a distributive lattice as well. Even more, the definition added to an ortholattice makes it orthomodular for quantum implications and distributive for the classical one. Based on this result a quantum implication algebra with a single primitive—and in this sense unique—implication is formulated. A corresponding classical implication algebra is also formulated. The algebras are shown to be special cases of a universal implication algebra.

1. INTRODUCTION

It is well known that there are five operations of implication in an orthomodular lattice which all reduce to the classical implication in a distributive lattice. (Kalmbach, 1983). It was therefore believed that implication algebras for these implications must all be different and such different algebras have explicitly been defined in the literature (Clark, 1973; Piziak, 1974; Abbott, 1976; Georgacarakos, 1980; Hardegree, 1981a). The systems were restrictive and uniquely determined a particular implication in an orthomodular lattice model.

In this paper, we show in Section 3 that one can formulate an implication algebra which can be modeled by either a bare orthomodular or distributive lattice only *after* choosing the representation of implication by means of the lattice operations. We arrive at such a formulation of implication algebra by using a novel possibility, obtained in Section 2, of turning an ortholattice

¹Atominstitute of the Austrian Universities, A-1020 Vienna, Austria; e-mail: pavicic@ati.ac.at; Department of Mathematics, University of Zagreb, HR-10001 Zagreb, Croatia; e-mail: mpavicic@faust.irb.hr.

²19 Locke Lane, Lexington, Massachusetts 02173; e-mail: nm@alum.mit.edu.

into either an orthomodular lattice or a distributive lattice by defining the join in the same way by means of either quantum or classical implications, respectively.

2. UNIFIED JOINS IN LATTICES

Definition 2.1. An ortholattice is an algebra $OL = \langle L^\circ, \perp, \cup \rangle$ in which the following conditions are satisfied for any $a, b, c, \in L^\circ$:

- L1.** $a \leq a^{\perp\perp} \ \& \ a^{\perp\perp} \leq a$
- L2.** $a \leq a \cup b \ \& \ b \leq a \cup b$
- L3.** $a \leq b \ \& \ b \leq a \Rightarrow a = b$
- L4.** $a \leq 1$
- L5.** $a \leq b \Rightarrow b^\perp \leq a^\perp$
- L6.** $a \leq b \ \& \ b \leq c \Rightarrow a \leq c$
- L7.** $a \leq c \ \& \ b \leq c \Rightarrow a \cup b \leq c$

where

$$a \leq b \stackrel{\text{def}}{=} a \cup b = b, \quad 1 \stackrel{\text{def}}{=} a \cup a^\perp$$

Also

$$a \cap b \stackrel{\text{def}}{=} (a^\perp \cup b^\perp)^\perp, \quad 0 \stackrel{\text{def}}{=} a \cap a^\perp$$

An ortholattice is orthomodular (OML) if the following conditions are satisfied for any $a, b \in L^\circ$.

- L8.** $a \rightarrow_i b = 1 \Rightarrow a \leq b \quad (i = 1, \dots, 5)$

and an ortholattice is distributive (DL) if the following condition is satisfied for any $a, b \in L^\circ$ (Pavičić, 1987)

- L9.** $a \rightarrow_0 b = 1 \Rightarrow a \leq b$

where the implications $a \rightarrow_i b$ ($i = 0, \dots, 5$) are defined as follows:

$$a \rightarrow_0 b \stackrel{\text{def}}{=} a^\perp \cup b \quad (\text{classical})$$

$$a \rightarrow_1 b \stackrel{\text{def}}{=} a^\perp \cup (a \cap b) \quad (\text{Sasaki})$$

$$a \rightarrow_2 b \stackrel{\text{def}}{=} b \cup (a^\perp \cap b^\perp) \quad (\text{Dishkant})$$

$$a \rightarrow_3 b \stackrel{\text{def}}{=} ((a^\perp \cap b) \cup (a^\perp \cap b^\perp)) \cup (a \cap (a^\perp \cup b)) \quad (\text{Kalmbach})$$

$$a \rightarrow_4 b \stackrel{\text{def}}{=} ((a \cap b) \cup (a^\perp \cap b)) \cup ((a^\perp \cup b) \cap b^\perp) \quad (\text{non-tollens})$$

$$a \rightarrow_5 b \stackrel{\text{def}}{=} ((a \cap b) \cup (a^\perp \cap b)) \cup (a^\perp \cap b^\perp) \quad (\text{relevance})$$

Theorem 2.1. (i) The equation

$$\mathbf{UJ}(i). \quad a \cup b = (a \rightarrow_i b) \rightarrow_i (((a \rightarrow_i b) \rightarrow_i (b \rightarrow_i a)) \rightarrow_i a)$$

is true in all orthomodular lattices for $i = 1, \dots, 5$ and in all distributive lattices for $i = 0$; (ii) an ortholattice in which $\mathbf{UJ}(i)$ holds is an orthomodular lattice for $i = 1, \dots, 5$ and a distributive lattice for $i = 0$.

Proof. (i) For $i = 1, \dots, 5$, the proofs are tedious, but straightforward expansions of the definitions, using the Foulis–Holland theorems extensively throughout. The following equations summarize the important intermediate results:

- UJ(3).1a.** $(a \rightarrow_3 b)^\perp \cap (b \rightarrow_3 a)^\perp = 0$
- UJ(i).1a.** $(a \rightarrow_i b) \cap (b \rightarrow_i a) = (a \cap b) \cup (a^\perp \cap b^\perp), \quad i = 4, 5$
- UJ(i).1b.** $(a \rightarrow_i b)^\perp \cap (b \rightarrow_i a) = a \cap b^\perp, \quad i = 3, 4, 5$
- UJ(i).1c'.** $(a \rightarrow_i b)^\perp \cup (b \rightarrow_i a) = a \cup b^\perp, \quad i = 3, 4$
- UJ(3).1c.** $(a \rightarrow_3 b) \cap ((a \rightarrow_3 b)^\perp \cup (b \rightarrow_3 a)) = (a^\perp \cap b^\perp) \cup (a \cap (a^\perp \cup b))$
- UJ(4).1c.** $((a \rightarrow_4 b)^\perp \cup (b \rightarrow_4 a)) \cap (b \rightarrow_4 a)^\perp = (a^\perp \cup b^\perp) \cap (a^\perp \cup b) \cap a$
- UJ(5).1c.** $(a \rightarrow_5 b)^\perp \cap (b \rightarrow_5 a)^\perp = (a \cup b) \cap (a \cup b^\perp) \cap (a^\perp \cup b) \cap (a^\perp \cup b^\perp)$
- UJ(i).1.** $(a \rightarrow_i b) \rightarrow_i (b \rightarrow_i a) = a \cup (a^\perp \cap b^\perp) \quad i = 1, 2, 3, 4$
- UJ(5).1.** $(a \rightarrow_5 b) \rightarrow_5 (b \rightarrow_5 a) = a \cup b^\perp$
- UJ(i).2.** $(a \cup (a^\perp \cap b^\perp)) \rightarrow_i a = a \cup b, \quad i = 1, 2, 3, 4$
- UJ(5).2.** $(a \cup b^\perp) \rightarrow_5 a = a \cup (a^\perp \cap b)$
- UJ(3).3a.** $(a \rightarrow_3 b)^\perp \cap (a \cup b)^\perp = 0$
- UJ(5).3a.** $(a \rightarrow_5 b) \cap (a \cup (a^\perp \cap b)) = (a \cap b) \cap (a^\perp \cap b)$
- UJ(i).3b.** $(a \rightarrow_i b)^\perp \cap (a \cup b) = (a \rightarrow_i b)^\perp, \quad i = 3, 4$
- UJ(5).3b.** $(a \rightarrow_5 b)^\perp \cap (a \cup (a^\perp \cap b)) = a \cap (a^\perp \cup b^\perp)$
- UJ(i).3c'.** $(a \rightarrow_i b)^\perp \cup (a \cup b) = a \cup b, \quad i = 3, 4$
- UJ(3).3c.** $(a \rightarrow_3 b) \cap ((a \rightarrow_3 b)^\perp \cup (a \cup b)) = (a^\perp \cap b) \cup (a \cap (a^\perp \cup b))$
- UJ(5).3c.** $(a \rightarrow_5 b)^\perp \cap (a \cup (a^\perp \cap b))^\perp = (a \cup b) \cap (a \cup b^\perp) \cap a^\perp$
- UJ(i).3.** $(a \rightarrow_i b) \rightarrow_i (a \cup b) = a \cup b, \quad i = 1, 2, 3, 4$
- UJ(5).3.** $(a \rightarrow_5 b) \rightarrow_5 (a \cup (a^\perp \cap b)) = a \cup b$

For $i = 3, 4$: $\mathbf{UJ}(i).1c$ follows from $\mathbf{UJ}(i).1c'$. For $i = 3, 4, 5$: $\mathbf{UJ}(i).1$ follows from $\mathbf{UJ}(i).1a,b,c$ and the definition of \rightarrow_i . For $i = 3$: $\mathbf{UJ}(3).3c$ follows

from UJ(3).3c'. For $i = 3, 5$: UJ(i).3 follows from UJ(i).3a,b,c and the definition of \rightarrow_i . For $i = 4$: UJ(4).3 follows from UJ(4).3b,c' and the definition of \rightarrow_4 . For $i = 1, \dots, 5$: UJ(i) follows from UJ(i).1, 2, 3.

For $i = 0$: The proof follows in a trivial way from the distributivity property and we omit it.

(ii) The nonorthomodular ortholattice O6 (Fig. 1a) is violated by UJ(i) for $i = 1, \dots, 5$. Each UJ(i) is therefore equivalent to the orthomodular law by Theorem 2(iii) of Kalmbach (1983, p. 22).

Nondistributive OM₆ (Fig. 1b) (Abbott, 1976) is violated by UJ(0). One easily shows that UJ(0) is equivalent to the distributive law. ■

Theorem 2.2. The equation

$$\mathbf{UO}(i). \quad a^\perp = a \rightarrow_i 0$$

is true in all orthomodular lattices for $i = 1, \dots, 5$ and in all distributive lattices for $i = 0$.

Proof. The proof is straightforward and we omit it. ■

3. QUANTUM AND CLASSICAL IMPLICATION ALGEBRAS

Many authors have given various systems of implication algebras. For example, systems for the Sasaki implication were formulated by Clark (1973), Piziak (1974), and Hardegree (1981a, b), for Dishkant's by Abbott (1976) and Georgacarakos (1980), and for the relevance implication by Georgacarakos (1980). In this section we give a system which is formulated by means of a primitive implication as the only operation. In doing so, in formulating definitions and theorems, we shall mostly follow Piziak (1974), Abbott (1976), and Georgacarakos (1980).

Definition 3.1. Let P be a nonempty set. We designate an element in P and call it 0. Let $\rightarrow: P \times P \rightarrow P$. Let $1 = 0 \xrightarrow{\text{def}} 0$. Then the triple $\mathcal{U}\mathcal{A} = \langle P, 0, \rightarrow \rangle$ is called a *universal implication algebra* provided the following



Fig. 1. (a) Ortholattice O6, (b) orthomodular lattice OM₆.

axioms and rules of inference R1–R4 are satisfied for all $a, b, c \in P$. The triple $\mathcal{Q}\mathcal{A} = \langle P, 0, \rightarrow \rangle$ is called a *quantum implication algebra* provided the following axioms and rules of inference R1–R5q are satisfied for all $a, b, c \in P$, and a *classical implication algebra* $\mathcal{C}\mathcal{A} = \langle P, 0, \rightarrow \rangle$ provided A1–A4, R1–R4, and R5c are satisfied for all $a, b, c \in P$.

Axioms.

- A1.** $a \rightarrow ((a \rightarrow 0) \rightarrow 0) = 1$ & $((a \rightarrow 0) \rightarrow 0) \rightarrow a = 1$
A2. $a \rightarrow ((a \rightarrow b) \rightarrow (((a \rightarrow b) \rightarrow (b \rightarrow a)) \rightarrow a)) = 1$
A3. $b \rightarrow ((a \rightarrow b) \rightarrow (((a \rightarrow b) \rightarrow (b \rightarrow a)) \rightarrow a)) = 1$
A4. $a \rightarrow 1 = 1$ & $(a \rightarrow (a \rightarrow 0)) \rightarrow (((a \rightarrow (a \rightarrow 0)) \rightarrow ((a \rightarrow 0) \rightarrow a)) \rightarrow a) = 1$

Rules of Inference.

- R1.** $a \rightarrow b = 1 \Rightarrow (b \rightarrow 0) \rightarrow (a \rightarrow 0) = 1$
R2. $a \rightarrow b = 1$ & $b \rightarrow c = 1 \Rightarrow a \rightarrow c = 1$
R3. $a \rightarrow c = 1$ & $b \rightarrow c = 1 \Rightarrow ((a \rightarrow b) \rightarrow (((a \rightarrow b) \rightarrow (b \rightarrow a)) \rightarrow a)) \rightarrow c = 1$
R4. $a \rightarrow b = 1$ & $b \rightarrow a = 1 \Leftrightarrow a = b$
R5q. $(a \rightarrow (b \rightarrow 0)) \rightarrow (((a \rightarrow (b \rightarrow 0)) \rightarrow ((b \rightarrow 0) \rightarrow a)) \rightarrow a) = 1$ & $a \rightarrow b = 1 \Rightarrow b \rightarrow a = 1$
R5c. $(a \rightarrow (b \rightarrow 0)) \rightarrow (((a \rightarrow (b \rightarrow 0)) \rightarrow ((b \rightarrow 0) \rightarrow a)) \rightarrow a) = 1 \Rightarrow b \rightarrow a = 1$

Theorem 3.1. Let $\mathcal{L} = \langle P, \perp, \cup \rangle$ be an orthomodular lattice OML. Define in \mathcal{L} the following operations:

- D1.** $1 \stackrel{\text{def}}{=} a \cup a^\perp$
D2. $0 \stackrel{\text{def}}{=} 1^\perp$
D3. $a \rightarrow b \stackrel{\text{def}}{=} a \rightarrow_i b$ where $i = \text{either } 1 \text{ or } 2 \dots \text{ or } 5$

Then the system $\mathcal{L}^{\mathcal{Q}\mathcal{A}} = \langle P, 0, \rightarrow \rangle$ is a quantum implication algebra $\mathcal{Q}\mathcal{A}$. The same is valid for a universal implication algebra $\mathcal{U}\mathcal{A}$.

Proof. The proof follows straightforwardly from Definition 2.1 and Theorem 2.1 and the following property of an orthomodular lattice (Pavičić, 1987):

- L10.** $a \cup b^\perp = 1$ & $a \leq b \Rightarrow b \leq a$ ■

Theorem 3.2. Let $\mathcal{L} = \langle P, \perp, \cup \rangle$ be a distributive lattice. Define in \mathcal{L} the following operations:

$$\mathbf{D1.} \quad 1 \stackrel{\text{def}}{=} a \cup a^\perp$$

$$\mathbf{D2.} \quad 0 \stackrel{\text{def}}{=} 1^\perp$$

$$\mathbf{D3.} \quad a \rightarrow b \stackrel{\text{def}}{=} a \rightarrow_0 b$$

Then the system $\mathcal{L}^{\mathcal{CA}} = \langle P, 0, \rightarrow \rangle$ is a classical implication algebra \mathcal{CA} . The same is valid for a universal implication algebra \mathcal{UA} .

Proof. The proof again follows straightforwardly from Definition 2.1 and Theorem 2.1 and the following well-known property of a distributive lattice:

$$\mathbf{L11.} \quad a \cup b^\perp = 1 \Rightarrow b \leq a \quad \blacksquare$$

Theorem 3.3. If $\mathcal{L} = \langle P, 0, \rightarrow \rangle$ is a quantum implication algebra \mathcal{QA} , then

$$\mathcal{L}^{\mathcal{QA}^*} = \langle P, \perp, \cup \rangle$$

is an orthomodular lattice OML, where \perp, \cup are defined in \mathcal{L} as follows:

$$\mathbf{D1.} \quad a^\perp \stackrel{\text{def}}{=} a \rightarrow 0$$

$$\mathbf{D2.} \quad a \cup b \stackrel{\text{def}}{=} (a \rightarrow b) \rightarrow (((a \rightarrow b) \rightarrow (b \rightarrow a)) \rightarrow a)$$

Moreover, $a \rightarrow b$ is determined as one of $a \rightarrow_i b, i = 1, \dots, 5$.

Proof. We first write all expressions of the form $a \rightarrow 0$ as a^\perp . Then we recognize all the $a \cup b$ expressions. Next, we write down all $a \rightarrow b = 1$ expressions as $a \leq b$ (which easily follows from R4). We are left with L1–L7 and L10, i.e., we obtain an orthomodular lattice. Hence, $a \rightarrow b$ from R4 must be one of $a \rightarrow_i b, i = 1, \dots, 5$, as given by L8. \blacksquare

Theorem 3.4. If $\mathcal{L} = \langle P, 0, \rightarrow \rangle$ is a classical implication algebra \mathcal{CA} , then

$$\mathcal{L}^{\mathcal{CA}^*} = \langle P, \perp, \cup \rangle$$

is a distributive lattice DL, where \perp, \cup are defined in \mathcal{L} as in Theorem 3.3. Moreover, $a \rightarrow b$ is determined as $a \rightarrow_0 b$.

Proof. We first write all expressions of the form $a \rightarrow 0$ as a^\perp . Then we recognize all the $a \cup b$ expressions. Next, we use R4 to write down all $a \rightarrow b = 1$ expressions as $a \leq b$. We are left with L1–L7 and L11, i.e., we obtain a distributive lattice. Hence, $a \rightarrow b$ from R4 must be $a \rightarrow_0 b$ as given by L9. \blacksquare

Theorem 3.5. If $\mathcal{L} = \langle P, 0, \rightarrow \rangle$ is a universal implication algebra $\mathcal{U}\mathcal{A}$, then

$$\mathcal{L}^{\mathcal{U}\mathcal{A}^*} = \langle P, \perp, \cup \rangle$$

is a distributive (an orthomodular) lattice, where \perp, \cup are defined \mathcal{L} as follows:

$$\mathbf{D1.} \quad a^\perp \stackrel{\text{def}}{=} a \rightarrow 0$$

$$\mathbf{D2.} \quad a \cup b \stackrel{\text{def}}{=} (a \rightarrow b) \rightarrow (((a \rightarrow b) \rightarrow (b \rightarrow a)) \rightarrow a)$$

and where \rightarrow maps into \rightarrow_i in the following way:

$$a \rightarrow_i b = a \rightarrow b; \quad i = 0 \quad \text{distributive} \quad (i = 1, \dots, 5 \quad \text{orthomodular})$$

Proof. We first write all expressions of the form $a \rightarrow 0$ as a^\perp . Then we recognize all the $a \cup b$ expressions. Next, we use R4 to write down all $a \rightarrow b = 1$ expressions as $a \leq b$. We are left with L1–L7, i.e., we obtain an ortholattice. Then, depending on whether we choose $a \rightarrow b$ from R4 to be either $a \rightarrow_0 b$ or $a \rightarrow_i b, i = 1, \dots, 5$, we obtain either a distributive or an orthomodular lattice. ■

4. CONCLUSION

In Section 2 we showed that an ortholattice, when the axiom UJ(i) defined in Theorem 2.1 is added to it, turns into an orthomodular lattice for $i = 1, \dots, 5$, i.e., for the so-called quantum implications, and into a distributive one for $i = 0$, i.e., for the classical implication. The axiom UJ expresses join by means of all these implications in a formally identical way.

In Section 3, in Definition 3.1 we employed the so expressed join to formulate quantum and classical implication algebras $\mathcal{Q}\mathcal{A}$ and $\mathcal{C}\mathcal{A}$, respectively. We have chosen the axiomatization from (Pavičić, 1987) because each expression of its axioms and rules of inference contains \cup and \perp at most once, and introduced join \cup and orthocomplementation \perp expressed by means of a primitive implication following UJ and UO from Theorems 2.1 and 2.2, respectively. In that way we obtain a quantum implication algebra which uses a single primitive implication. Hence, the main difference between our and all previous quantum implication algebras is that each of the latter corresponds to a particular quantum implication (one of five), while ours does not. (Besides, to our knowledge, only for three of five implications, implication algebras have been formulated so far.) As soon as we define the lattice operations on the algebra, by means of D1–D2 from Theorem 3.3, we open the orthomodular lattice possibility to express the primitive implication on the algebra by means of any of the five quantum implications, but this is

exactly the sense in which the obtained quantum implication algebra $\mathcal{Q}\mathcal{A}$ generalizes not only orthomodular lattices, but also all previously obtained quantum implication algebras.

As also shown in Section 3, the two generalized quantum and classical implication algebras can themselves be generalized by dropping the rules R5 from Definition 3.1. The obtained universal implication algebra $\mathcal{U}\mathcal{A}$ (see Theorem 3.5) contains algebras $\mathcal{Q}\mathcal{A}$ and $\mathcal{C}\mathcal{A}$, depending on how we choose to represent the implication. It would be interesting to investigate what other systems the algebra $\mathcal{U}\mathcal{A}$ can give under different definitions of the lattice operations.

NOTE ADDED IN PROOFS

A step by step proof of Theorem 2.1 the reader can find at the following address: www1.shore.net/~ndm/java/augl/mmxplorer.html

ACKNOWLEDGMENTS

M.P. acknowledges supports of the Ministry of Science of Croatia and the Technical University of Vienna, Austria. He would also like to thank Johann Summhammer, Atominstute of Austrian Universities, Vienna, for his hospitality.

REFERENCES

- Abbott, J. C. (1976). Orthoimplication algebra, *Studia Logica*, **35**, 173–177.
- Clark, I. D. (1973). An axiomatisation of quantum logic, *Journal of Symbolic Logic*, **38**, 389–392.
- Georgacarakos, G. N. (1980). Equationally definable implication algebras for orthomodular lattices, *Studia Logica*, **39**, 5–18.
- Hardegree, G. M. (1981a). Quasi-implication algebras, Part I: Elementary theory, *Algebra Universalis*, **12**, 30–47.
- Hardegree, G. M. (1981b). Quasi-implication algebras, Part II: Structure theory, *Algebra Universalis*, **12**, 48–65.
- Kalmbach, G. (1983). *Orthomodular Lattices*, Academic Press, London.
- Pavičić, M. (1987). Minimal quantum logic with merged implications, *International Journal of Theoretical Physics*, **26**, 845–852.
- Piziak, R. (1974). Orthomodular lattices as implication algebras, *Journal of Philosophical Logic*, **3**, 413–438.